On the Probabilities of Branch Durations and Stratigraphic Gaps in Phylogenies of Fossil Taxa when Rates of Diversification Vary over Time

Peter J. Wagner

Dept. of Earth & Atmospheric Sciences, and School of Biological Sciences

316 Bessey Hall

University of Nebraska, Lincoln USA

pjwagner3@gmail.com

Abstract

The time separating the first appearances of species from their divergences from related taxa affects assessments of macroevolutionary hypotheses about rates of anatomical or ecological change. Branch durations necessarily posit stratigraphic gaps in sampling within a clade over which we have failed to sample predecessors (ancestors) and over which there are no divergences leading to sampled relatives (sister-taxa). The former reflects only sampling rates whereas the latter reflects sampling, origination and extinction rates. Because all three rates vary over time, the probability of a branch duration of any particular length will differ depending on when in the Phanerozoic that branch duration spans. Here, I present a birth-death-sampling model allowing interval-to-interval variation in diversification and sampling rates. Increasing either origination or sampling rates increases the probability of finding sister taxa that diverge both during and before intervals of high sampling/origination. Conversely, elevated extinction reduces the probability of divergences from sampled sister-taxa before and during intervals of

elevated extinction. In the case of total extinction, a Signor-Lipps will reduce expected sister-taxa leading up to the extinction, with the possible effect stretching back many millions of years when sampling is low. Simulations indicate that this approach provides reasonable estimates of branch duration probabilities under a variety of circumstances. Because our current probability models for describing morphological evolution are not as advanced as our methods for inferring diversification and sampling rates, branch duration priors allowing for time-varying diversification and could be a potent tool for phylogenetic inference with fossil data.

Resubmitted to *Paleobiology* on 2018-06-14.

Introduction

The divergence times implicit to phylogenetic relationships among fossil taxa almost always posit durations over which we have failed to sample taxa from a clade and thus imply gaps in stratigraphic sampling (Smith 1988). Given the challenges that easily fossilized characters present to phylogenetic analyses, such as abundant homoplasy and correlated change (e.g., Wagner 2000a; Smith 2001a; Wagner and Estabrook 2015; Wright et al. 2016; Sansom and Wills 2017), the plausibility of the stratigraphic gaps implicit to branch durations is critical for evaluating the plausibility of general sets of relationships (Fisher 1994). Within the context of particular cladistic relationships, the probabilities of branch durations are important for assessing macroevolutionary hypotheses about consistency of rates of anatomical change and/or ecological change (Ruta et al. 2006, Friedman 2010, Bapst et al. 2016, Halliday and Goswami 2016) and for accommodating the effects of rate variation and even character coding practices on methods that try to calibrate divergence times using local morphological “clocks” (i.e., tip-dating; Ronquist et al. 2012; Matzke and Wright 2016; Matzke and Irmis 2018).

The probabilities of branch durations and associated stratigraphic gaps reflect not just sampling rates, but also origination and extinction rates (e.g., Foote et al. 1999; Stadler 2010; Bapst 2013; Heath et al. 2014). Ideally, if we can put prior probabilities on divergence times given extrinsic information about these rates, then analyses using birth-death-sampling models to estimate divergences times and basic phylogenetic relationships among fossil taxa (e.g., Wright 2017; Wright and Toom 2017) will be more robust to local idiosyncrasies in rates or modes of morphological change. Marshall (2017) draws attention to the fact that existing birth-death-sampling models for estimating prior probabilities of phylogenetic branches do not

take into account existing paleobiological estimates of rates of diversification or sampling. Although some models do allow for heterogeneity in these rates among taxa (e.g., Stadler et al. 2013, Stadler et al. 2018), existing methods do not accommodate the clade-wide shifts in these rates from one interval to the next that the fossil record demonstrates. Currently, our methods for estimating diversification and sampling are more advanced than are our models for describing morphological evolution on which phylogenetic and divergence time inferences rely; thus, flexible birth-death-sampling models that can generate prior probabilities for phylogenies using temporal variation in diversification and sampling could improve the accuracy of estimated phylogenies and divergence times for analyses of fossil taxa.

The basic problem and a general model

The problem that we need to address is, what is the probability that two sampled species could have a posited divergence time without one or the other having had a more recent divergence time with another sampled taxon? Every sampled species represents the tip of a lineage extending back to the origin of the clade being studied. Barring exceptional circumstances such as species-flocking (e.g., Yacobucci 1999), the lineages leading from the base of the clade to most species are united as single lineages that diverged at some point in the clade's history. Thus, the *branch duration* connecting any sampled Species X to its *divergence time* from the rest of the clade usually will span less time than the extended lineage leading to the base of the clade. (I use “branch duration” rather than “branch length” because the latter term also is used for expected character change along a phylogenetic branch; e.g., Lewis 2001.) Similarly, the branch duration separating the node linking Species X to other sampled species usually will span less time than the remainder of that extended lineage leading to the base of the clade.

79

80 Consider two hypothetical taxa, Species A and Species I (Fig. 1). Species A appears 304 Ma and
 81 Species I appears 306 Ma. The particular phylogeny we are evaluating posits that the two species
 82 diverged from a common ancestor (here represented by Species D) at 310 Ma. The maximum
 83 branch durations preceding either species are 6-myr for Species A and 4-myr for Species I (i.e.,
 84 the dark gray [dark blue online] bars in Fig. 1A). One evaluation of this divergence time
 85 estimate is the probability of zero finds given the branch durations (Wagner 1995a, 2000b;

Huelsenbeck and Rannala 1997). This assesses the probability that we failed to sample an immediate predecessor of Species A and Species I. However, Foote et al. (1999) make the important point that unless sampling levels are very high, most divergence times are set by sampling sister taxa rather than predecessors. Thus, if we sample any of the light gray [yellow online] lineages, then we will have a more recent divergence time and shorter branch duration for either Species A or Species I. For example, if we eventually sample Species F and correctly reconstruct the phylogeny, then we will reconstruct a divergence between Species A and Species F at 307 Ma. The branch duration leading from Species A to the rest of the clade reduces to 3-myr; the remaining 3-myr linking Clade A+F to the base of the clade now is the branch duration preceding the clade including Clade A+F. Thus, estimating the probability of a branch duration and divergence time requires estimating the probability of no sampled predecessors and the probability of no additional divergences leading to sampled sister taxa.

As Bapst (2013) emphasizes, the probability of any divergence time (and its implicit branch duration and stratigraphic gap within a phylogeny) reflects three rates: origination, extinction and sampling rates all affect the probability of a divergence time. Origination and extinction both have two effects on this probability. Origination rates provide the expected number of cladogenetic events along any branch leading to the base of the tree. Both origination and extinction affect the expected number of progeny that any one cladogenetic event will ultimately generate: as origination increases and/or extinction decreases (i.e., as net-diversification increases), we expect more taxa to evolve that could become sampled sister taxa. That in turn reduces the probability of deep divergences by increasing the probability of finding at least one descendant of a divergence even if sampling rates themselves do not increase. Decreasing

extinction has a similar effect, as increasing the durations of individual species increases the probability that at least one descendant of a divergence is found even if neither origination nor sampling increase (Foote 1996). Figure 1 illustrates how diversification affects expected branch durations and divergence times. There are three branching events leading to Species A compared to only two branching events leading to Species I; thus, all else being equal, it is more probable that subsequent sampling will add a sister-taxon to Species A rather than to Species I. Moreover, one of Species A's possible sisters, the C+E+F clade, offers many more opportunities for sampling than do any of the other divergences. All else being equal, the probability of sampling some descendant of that "C" divergence is greater than the probability of sampling descendants of any other divergence.

These expectations are independent of particular speciation models. Re-rendering Fig. 1A as either budding (Fig. 1B) or bifurcating (Fig. 1C) speciation models leaves the same branch durations leading from Species A and Species I to the base of the tree, and the same sum of durations for additional potential sister-taxa. The difference is that "predecessors" and "ancestors" are basically identical given bifurcation because ancestral morphotypes anagenetically change at cladogenesis, whereas predecessors include only those members of an ancestral species extant prior to cladogenesis under the budding model. (Under both models, the earliest members of Species A and Species I also are "predecessors.") Under the latter model, the probability of sampling an ancestor is greater than the probability of sampling a predecessor (Foote 1996). To avoid implying that ancestors necessarily become pseudoextinct at speciation, I will use the term "predecessor" for the analogs of the blue bars in Fig. 1 throughout this paper.

Bapst (2013) and Didier et al (2017) extend the work of Foote et al. (1999) by deriving the probability of sampling a species or any successors (i.e., a clade of unknown size) given time-homogeneous origination, extinction and sampling. These birth-death-sampling models (= fossilized-birth-death sampling sensu Heath et al. 2014; Gavryushkina et al. 2016; Zhang et al. 2016; Stadler et al. 2018) offer great potential as tools for assessing the branch durations and divergence times implicit to rival phylogenetic hypotheses, and thus for assessing different models of anatomical evolution favoring those rival phylogenetic hypotheses. Moreover, because we have abundant paleobiological methods for estimating origination, extinction and sampling from the fossil record, this represents a relatively rare case where we can empirically justify the rate parameters that we need to estimate prior probabilities of branch durations and divergence times in Bayesian analyses.

The same paleobiological methods that provide us with diversification and sampling rates also indicate that all three rates all vary over time (e.g., Connolly and Miller 2001; Foote 2001, 2003; Liow and Finarelli 2014; Alroy 2014; Finarelli and Liow 2016). What we require to accommodate interval-to-interval variation in origination, extinction and sampling is a general model that will estimate: 1) the probability of a species appearing in some interval generating any sampled specimens either of itself or some descendant species given the diversification and sampling rates of that interval and when in that interval the first species originated; 2) the probabilities that the first species sends $1 \dots \infty$ successors (with the successors possibly including the first species itself) into the next interval given the origination and extinction rates of that first interval; and, 3) the probability that any of those successors generate at least one sampled species given the origination, extinction and sampling rates of subsequent intervals.

The probability of sampling a species or any successors from any one interval

In Fig. 1, we posit that Species A and Species I are each other's closest sampled relative. The divergence time demands a branch duration and stratigraphic gap over which we failed to sample any specimens and over which no other sampled relatives diverged. If diversification and sampling rates are constant, then the expected number of sampled species or clades that should be more closely related to Species A than is Species I is $\Phi\lambda d$, where λ is the origination rate, Φ is the proportion of branching events for which any successors are sampled (i.e., the probability of sampling a clade of unknown size sensu Bapst 2013), and d is the branch duration leading to Species A (Table 1). Φ reflects sampling (ψ) and the summed species durations (S) expected given λ and μ (Foote et al. 1999). For example, the C'+E+F clade (Fig. 1A) has $S=7.75$ lineage-million-years (Lmyr). Bapst (2013 eqn. 2) extends this logic to estimate the probability of sampling a species or any successors (i.e., a clade of unknown size) given constant λ , μ and ψ as:

$$\Phi = 1 - \sum_{K=1}^{\infty} \frac{\lambda^{(K-1)} \mu^K \binom{2K-2}{K-1}}{K(\lambda + \mu + \psi)^{(2K-1)}} \quad (1a)$$

where $K=1 \dots \infty$ represents the possible richness generated by any one divergence. Didier et al (2017; Appendix 5; see also King in Bapst 2016) estimate Φ using a quadratic:

$$\Phi = 1 - \frac{(\lambda + \mu + \psi) - \sqrt{(\lambda + \mu + \psi)^2 - 4\lambda\mu}}{2\lambda} \quad (1b)$$

176

Table 1. Glossary.	
d	Branch duration, i.e., the time (and chronostratigraphic gap) separating the first appearance of a species (or the splitting time of a node) from the divergence linking it to the rest of the clade.
t	Time elapsed after a divergence over which a clade of any size might accrue new species, sampled specimens or become extinct.
T	Duration of interval with unique diversification and/or sampling rates.
S	Sum of species durations within a clade, usually given in lineage-million-years (Lmyr).
λ_i	Cladogenesis (origination) rate in interval-slice i .
μ_i	Extinction rate in interval-slice i .
ψ_i	Sampling rate in interval-slice i .
Φ_i	Probability of ever sampling a species and/or any successors (= a clade of unknown size) when only one species is present at the outset of time-slice i .
Φ'_i	Probability of ever sampling a species and/or any successors within a particular interval when only one species is present at the outset of interval-slice i .
v_i	Probability of sampling a species and/or any successors <i>within</i> interval-slice i .
ζ_i	Probability of sampling a species and/or any successors <i>after</i> interval-slice i when only one species is present at the outset of time-slice i .
$E[d_i]$	Expected branch duration for a species with first appearance in interval-slice i given origination, extinction and sampling rates. This is also the expectation for an unsampled ancestor with descendants that diverged in interval-slice i .

177 Bapst's and Didier et al.'s equations generate very similar estimates of Φ , and in both cases the
 178 probability of sampling a species or any successors increases as:

- 179 1. λ increases (increasing S by generating more species);
- 180 2. μ decreases (increasing S by generating longer-lived species);
- 181 3. ψ increase (increasing the probability of sampling per Lmyr).

182

183 To estimate Φ given shifting diversification and/or sampling, we first need to estimate Φ' , the
 184 probability of sampling a clade of unknown size over some interval i of duration T_i with its rates
 185 of λ_i , μ_i and ψ_i . (Note that time now refers to time elapsed after a divergence, not time prior to a
 186 first appearance or the splitting from a common predecessor.) First, we calculate the probability
 187 that one lineage will result in $K = 0 \dots \infty$ lineages at any time from $t = 0 \dots T$ (Fig. 2). When $\lambda \neq \mu$:

$$188 \quad P[K = 0 | \lambda, \mu, t] = \frac{\mu \times (e^{(\lambda-\mu)t} - 1)}{(\lambda \times e^{(\lambda-\mu)t} - \mu)} \quad (2)$$

189 (Raup 1985: eqn. A13) and

190
$$P[K \geq 1 | \lambda, \mu, t] = \left(1 - \frac{\mu \times (e^{(\lambda-\mu)t} - 1)}{(\lambda \times e^{(\lambda-\mu)t} - \mu)}\right) \times \left(1 - \frac{\lambda \times \mu \times (e^{(\lambda-\mu)t} - 1)}{\mu \times ([\lambda \times e^{(\lambda-\mu)t}] - \mu)}\right) \times$$

191
$$\left(\frac{\lambda \times \mu \times (e^{(\lambda-\mu)t} - 1)}{\mu \times ([\lambda \times e^{(\lambda-\mu)t}] - \mu)}\right)^{K-1} \quad (3)$$

192 (Raup 1985: eqn. A13, A17). When $\lambda=\mu$:

193
$$P[K = 0 | \lambda, \mu, t] = \frac{\lambda t}{1 + \lambda t} \quad (4)$$

194 (Raup 1985: eqn.. A12), and

195
$$P[K \geq 1 | \lambda, \mu, t] = \frac{(\lambda t)^{K-1}}{(1 + \lambda t)^{K+1}} \quad (5)$$

(Raup 1985: eqn. A14). Fig. 2 illustrates these in 9 combinations of rates over some interval with $T=2.5$. Both λ and μ give expectations over $t=1$ (e.g., rates per million years for an interval of 2.5 million years).

Integrating over the individual probability curves for K successors of one species gives the $E[t_K]$, the expected amount of time within interval i in which K successors of the original species are present (Fig. 3). We can use this to estimate $E[S]$, the expected sum of durations within an interval:

$$E[S] = \sum_{K=1}^{\infty} (K \times E[t_K])$$

(Eq. 6)

i.e., the number of lineages (K) times the expected time with that number of lineages ($E[t_K]$) summed over all possible K . (Because $E[t_K]$ rapidly declines to very low numbers, the summation can be quickly truncated.) If we have Interval A with $\lambda=\mu=0.4 \text{ myr}^{-1}$ (Fig. 3A), then:

$$\begin{aligned} E[S_A] &= (1 \times 1.250) + (2 \times 0.625) + (3 \times 0.313) + (3 \times 0.156) + (4 \times 0.078) \dots \\ &= 2.5 \text{ Lmyr.} \end{aligned}$$

If we have Interval B with $\lambda=0.566$ and $\mu=0.4$ (Fig. 3B), then:

$$\begin{aligned} E[S_B] &= (1 \times 1.125) + (2 \times 0.716) + (3 \times 0.456) + (3 \times 0.290) + (4 \times 0.185) \dots \\ &= 3.1 \text{ Lmyr.} \end{aligned}$$

A first approximation of the probability of sampling a lineage present at the outset of any interval X or any successors to that species is:

$$\Phi'_X = 1 - e^{-\psi_X E[S_X]}$$

(Eq. 7).

Therefore, if $\psi_A < 1.24 \times \psi_B$, then we expect to sample more clades diverging at the onset of Interval B than we would for Interval A.

Simulations show that Eq. 7 overestimates Φ' . At $\lambda=\mu=\psi=0.4$ and $T=2.5$, equations 6+7 estimate $\Phi'=0.632$, whereas simulations starting using these same parameters generate sampled species in only 53.3% of runs (see also Supplementary Fig. S1). The culprit is the failure of Eq. 6 to account for total extinction within the interval eliminating any possibility of additional sampling regardless of the remaining expected S (see Supplementary Fig. S2). One solution is to divide intervals into m interval-slices so that each per-slice ψ is low and then estimate: 1) v_i , the probability of a species extant at the outset having sampled members or successor in interval-slice i (Eq. 7); and, 2) ζ_i , the probability of a lineage present at the outset of interval-slice i having sampled successors in interval-slices $i+1$ to m . The former parameter, v , is the probability of finding a species and/or any successors in an interval-slice:

$$v_i = 1 - e^{-\psi_i E[S_i] t_i}$$

(Eq. 7a)

233 where ψ_i is the sampling rate for the interval, S_i is the expected total progeny in the interval-slice,
 234 and t_i is duration of the interval-slice. The latter parameter, ζ , is the probability of $K=1 \dots \infty$ total
 235 progeny yielding at least one sampled species times given “future” λ , μ and ψ multiplied by the
 236 probability of the initial species having $K=1 \dots \infty$ successors (again, possibly including the
 237 original species) at the end of the interval-slice i given λ_i and μ_i (e.g., Fig. 5):

$$\zeta_i = \sum_{K=1}^{\infty} (P[\geq 1 \text{ Finds } |K, \Phi'_{i+1}] \times P[K \text{ Successors } |\lambda_i, \mu_i])$$

(Eq. 8).

For the final interval-slice m , $\zeta_m = 0$ because there are no opportunities to sample successors later in the interval. Thus, $\Phi'_m = v_m$ and we can solve $\zeta_{1 \dots (m-1)}$ recursively. The maximum possible value for ζ_i is the probability of any successors surviving interval-slice i : at $\lambda=\mu=0.4$ and $T_i=0.25$, the probability of extinction is 0.091 (Fig 4A) and the maximum possible ζ_i is 0.909. Now:

$$\Phi'_i \cong 1 - ([1 - v_i] \times [1 - \zeta_i])$$

(Eq. 9).

This explicitly accommodates the probability that a lineage present at the outset of an interval is extinct with no successors by (say) the 5th of $m=10$ interval-slices.

Simulations (Fig. 4; Supplementary Fig. S1) show that Eq. 9 rapidly converges to appropriate estimates of Φ'_1 at $m \geq 10$ over the range of diversification parameters used in Figs. 2-4. Thus, if we typically have intervals of (say) 2.5 million years over which we have estimates of λ , μ and ψ , then 100-kyr interval-slices should be adequate to estimate Φ'_1 .

Another useful aspect of calculating Φ'_i for multiple slices within an interval is that divergences could happen at any time within an interval if origination is continuous through time. Suppose that the precision of divergence dates that we will consider are in 100-kyr increments: that is, we will ask whether a divergence happened at 450.1 Ma, 450.0 Ma, 449.9 Ma, etc. If these divergences are within a 2.5-myr chronostratigraphic interval, then we then we need to know $\Phi'_{1 \dots 25}$. Φ'_i will decrease as i increases: we have both less time to accrue lineage durations and

less time in which to sample them. If the interval has $\lambda=\mu=\psi=0.4 \text{ myr}^{-1}$, then: $\Phi'_1=0.536$,
 $\Phi'_{13}=0.383$, and $\Phi'_{25}=0.039$. Thus, even if the probability of a divergence is the same at any
point within the interval, the probability of a divergence that generates a sampled species *within*
that interval declines over time.

The probability of sampling a species or successors over 2+ intervals with different diversification and/or sampling rates

The exercise above applies sampling within one interval. However, this is just the particular
case where all m diversification and sampling rates are the same and $v_1=v_2=\dots=v_m$. We can
calculate Eq. 9 just as easily using $v_1 \neq v_2 \neq \dots \neq v_m$, as would be the case if either of the diversification
rates or the sampling rate varied in every interval-slice. Similarly, ζ_i is just as easy to calculate if
 λ_i and/or μ_i differ in every interval-slice. What I will do here is intermediate: interval-slices from
different intervals will use particular values of λ , μ and ψ so that each v_i within an interval is
identical and each ζ_i within an interval uses the same values of λ and μ . The one difference is
that ζ_m will be greater than zero for all intervals save the last, and all ζ_i from those intervals will
reflect subsequent shifts in diversification and/or sampling.

Suppose that we are analyzing a clade of Ordovician gastropod species known through the
Katian. Suppose further that data from the Paleobiology Database coupled with an external
stratigraphic database allow us to estimate separate diversification and sampling rates for
gastropods as a whole for three divisions of the Katian, corresponding to Bergström et al.'s
(2009) Ka1, Ka2+Ka3 (hereafter: Ka2-3), and Ka4: $\lambda_{\text{Ka1}}=0.269$, $\lambda_{\text{Ka2-3}}=0.348$, $\lambda_{\text{Ka4}}=0.312$;

282 $\mu_{Ka1}=0.293$, $\mu_{Ka2-3}=0.495$, $\mu_{Ka4}=0.514$; and, $\psi_{Ka1}=0.147$, $\psi_{Ka2-3}=0.155$, $\psi_{Ka4}=0.122$ (Fig. 6).

283 These parameters come from capture-mark-recapture analysis (Connolly and Miller 2001; Liow

284 and Nichols 2010) using Paleobiology Database data, with the analysis modified to allow for

285 lognormal distributions of sampling rates (Wagner and Marcot 2013) and each ψ_{Ka} the median of

286 those lognormal distributions. Each 100-kyr interval-slice within any one interval will have the

287 same probability of a divergence in that interval-slice being sampled immediately:

288 $v_{i \cdot Ka1}=1.46 \times 10^{-2}$, $v_{i \cdot Ka2-3}=1.53 \times 10^{-2}$, $v_{i \cdot Ka4}=1.20 \times 10^{-2}$ (Fig. 6A). Each 100-kyr interval-slice

289 within an interval also shares the same probability of a species present at the outset having 0... ∞

290 successors at the outset of the next 100-kyr interval-slice (Supplementary Fig. S3). However, ζ_i

291 for a Ka2-3 divergence (i.e., the probability of sampling the species or successors in subsequent

interval-slices), now reflects diversification and sampling in both Ka2-3 and Ka4 (Fig. 6). This in turn means that we now have Φ_i for each interval-slice within all three intervals: i.e., the probability of ever sampling a descendant of a divergence occurring in that 100-kyr interval-slice at some point in the time-range encompassed by our study. Simulations using the same origination, extinction and sampling rates per interval indicate that equations 7a and 8 combined generate accurate estimates of Φ_i for all interval-slices (Supplementary Fig. S4). This in turn indicates that Eq. 6 is accurately estimating $E[S]$ (expected summed durations evolving from a divergence) for different interval-slices given heterogeneous λ and μ .

The probability of branch durations and divergence times given shifting rates of diversification and/or sampling

We now can estimate the probability that a branch duration leading to a sampled species (or to the basal divergence of a sampled clade) diverges from the rest of the clade in any particular interval-slice. Following the gastropod example above, suppose that we are analyzing a group of bellerophonitoids through the Katian. One of the species is *Salpingostoma richmondensis*, which is first known from rocks at the base of the *Aphelognathus divergens* conodont zone in Ka4. What is the probability that *S. richmondensis* diverged from its closest relative prior to the Katian? Based on Gradstein et al. (2012), we assign an earliest appearance date of 447 Ma for *S. richmondensis*, i.e., about halfway through Ka4. (I will disregard uncertainty in the time-scale here.) Given that the Katian starts at 453 Ma, we are positing a minimum branch duration of $d=6$ myr. The probability of *failing* to sample a predecessor of *S. richmondensis* over $d=6$ -myr is the probability of missing it in every interval-slice i :

$$e^{-\sum_{i=1}^{FA-1} \psi_i d_i}$$

(Eq. 10),

where FA-1 is the interval-slice preceding the first appearance (here, the interval-slice corresponding to 447.1 Ma), d_1 is the time encompassed by each interval-slice (here, 100-kyr), and ψ_i is the sampling rate of that interval. (Fig. 6B gives the compliment of this, i.e., the probability of sampling a predecessor per interval-slice). Given the sampling rates from above, this comes to $p=0.420$.

The probability of there being no divergences leading to a sister-taxon of *Salpingostoma richmondensis* sampled in the Katian reflects the three separate origination rates and Φ_i at any given point in time:

$$\prod_{i=1}^{FA-1} \sum_{n=1}^{\infty} (P[n \text{ branchings} | \lambda_i d_i] \times [1 - \Phi_i]^n)$$

(Eq. 11),

where $P[n \text{ branchings} | \lambda_i]$ is the Poisson probability of n cladogenetic events in an interval-slice given the origination rate of that interval and the duration of each interval-slice (here, 100-kyr), and $[1-\Phi_i]^n$ is the probability of n divergences yielding no sampled sister taxa. Here, this generates $p=0.574$. We usually will get almost identical results estimating the probability of no divergences leading to sampled sister taxa as:

$$e^{-\sum_{i=1}^{FA-1} \Phi_i \lambda_i d_i}$$

because the probability of just *one* branching event is nearly equal to the probability of there being *any* branching events. However, at origination rates comparable to those estimated for major radiations, the probability of two branching events over even short intervals is great

enough that not allowing for $n=2$ yields notable underestimates of the probability of a sampled sister taxon. (Note that even very high λ , we can truncate the summation by $n=5$.)

The compliment of Eq. 11 is the probability that any given interval-slice includes a divergence leading to a sampled sister-taxon of *Salpingostoma richmondensis* (Fig. 6B). This increases not just as Φ increases (Fig. 6A), but also as λ increases: thus, the probability of a divergence leading to a sampled sister-taxon of *S. richmondensis* peaks early in Ka2-3, despite the edge-effect of ending the study after the next interval (Fig. 6B). Because ψ_{Ka2-3} is only marginally higher than ψ_{Ka1} and ψ_{Ka4} is less than ψ_{Ka1} this necessarily is driven by the λ_{Ka2-3} being greater than λ_{Ka1} . The combination of elevated λ and elevated ψ in Ka2-3 also means that these interval-slices have the highest probabilities of providing *S. richmondensis*' divergence from either a sampled predecessor or a sampled sister-taxon.

We now calculate the probability of branch duration d as the probability of no sampled predecessors (i.e., one minus eq. 10) times the probability of no divergences leading to sampled sister-taxa (i.e., one minus eq. 11). Thus, the probability that *Salpingostoma richmondensis* has a divergence time from other bellerophontoids no later than the base of the Katian given the shifting diversification and sampling rates of the Katian (and without the possibility of sampling post-Katian sister-taxa) is $0.420 \times 0.574 = 0.241$ (Fig. 6C). Finally, note that if we do not include any post-Katian species, then the expected branch duration for *S. richmondensis* ($E[d]$), i.e., the point where the probability of the branch duration is 0.5, is 3.1-myr.

Effects of simple shifts in sampling and/or diversification rates on expected stratigraphic gaps and branch durations

The specific effects of each parameter are difficult to appreciate when origination, extinction and sampling all vary from one interval to the next as in the Katian gastropod example. Moreover, the short span of time encompassed by the Katian data creates a major edge effect: Φ_i steadily declines throughout the Katian because no post-Katian descendants of Katian divergences can be sampled. Most studies include many more than one stage and thus might have many intervals without such an edge-effect. Indeed, just how long before the conclusion of a study (or before the total extinction of a clade) we need to go before this edge effect disappears is in itself a question of interest.

Therefore, I will present simple examples in which (with one exception) two of the rate parameters are held constant while one varies over time. This will emphasize how the particular parameters affect the probabilities of branch durations and divergence times. I will first cover shifts in sampling with constant diversification. I then will cover some simplified versions of major evolutionary events given constant sampling. The latter examples are particularly important, not simply because they emphasize the strong effects of both origination and extinction rates on expected branch durations and stratigraphic gaps within a phylogeny, but because paleobiologists often target species spanning such events when conducting phylogenetic analyses. In each example, I will use 6 intervals of 2.5-myr each, with estimates of Φ based on 100-kyr interval-slices. The “standard” diversification rates are set to 0.4 myr^{-1} so that we expect one event per interval. In all cases, I will use different median sampling levels that range from an expectation of one sample per interval to one sample per 100 intervals. To avoid the

edge-effect from the Katian gastropod example, all of the examples (save one) are followed by 12 intervals with “normal” λ , μ and ψ .

For each combination of parameters and parameter shifts, I present the following: 1) Φ , the probability of sampling a species or any successors (= a clade of unknown size); 2) the probability of a divergence leading to a sampled sister-taxon given expectation $\lambda_i \Phi_i$; 3) the probability of truncating a branch duration in interval-slice i either by sampling a predecessor or from the divergence of a subsequently sampled sister-taxon; 4) the expected branch duration ($E[d]$) given a sampled species with that first appears in interval-slice i . The last represents the branch duration where the probability of encountering either a sampled predecessor or the divergence time of a sampled sister-taxon reaches 0.5. Note that this $E[d]$ also applies to the expected branch duration leading to a node that diverges in interval-slice i .

Finally, for Φ over time, I also include the results from 10000 simulations (dots in Figs. 7A, 8). Each simulation begins with a single lineage, with both lifetimes and descendants per time-slice allowing for shifts in either diversification or sampling. Lifetimes and progeny of descendants are simulated the same way, albeit using rates specific to their lifetimes. Interval-specific rates of sampling are used when sampling changes. The results show the proportion of simulated lineages that generate 1+ sampled species.

Effects of shifts in sampling rates (ψ)

Although there are several evolutionary scenarios positing general shifts in origination and/or extinction rates, temporal patterns in sampling rates (ψ) do not have analogous simple models.

404 The biggest control on ψ is the amount of available sedimentary rock from the right
 405 environmental type (Raup 1972; Sheehan 1977; Peters and Foote 2001; Smith 2001b),
 406 which can vary fairly haphazardly over time. In the absence of any such templates, I use two
 407 shifts in ψ , where one interval has twice the median sampling rates and another has half the
 408 median sampling rates (Fig. 7). A key feature is that the effects of shifts in sampling on the
 409 probability of sampling a species or any descendants (Φ) *precedes* shifts in ψ (Fig. 7A):
 410 compared to species appearing in other intervals, species appearing shortly before elevated

Figure 7. Effects of variable sampling over time with constant origination and extinction ($\lambda=\mu=0.40$). Five median sampling rates (ψ) are illustrated, with ψ relative to the median given on by the gray bars (2nd y-axis). The six intervals of 2.5-myr are preceded by six others with median ψ . A). Φ , the probability of sampling a species or any successors given that the species is extant in an interval-slice. Yellow dots give expected Φ given 10000 simulations using ψ (and λ and μ) of the intervals of a species lifetime. B) The probability of a sampled sister-species diverging in each interval-slice. C) The probability of a phylogenetic divergence in each interval-slice from either sampling a predecessor or from the divergence of a subsequently sampled sister-taxon. D) Expected durations (d) for a species first appearing in each interval-slice. This also applies to a node that diverged in the same interval-slice. E[d] is the branch duration where P[no ancestors or divergences] reaches 0.50 (see Fig. 6C).

sampling have an higher probability of surviving to and/or having descendants in the interval of high sampling even if they do not have elevated probabilities of numerous descendants and/or long durations. Conversely, Φ is depressed prior to the fourth interval because species arising then have much lower probability of having descendants present during “normal” sampling intervals. We see the same patterns in reverse within the intervals of elevated and depressed ψ : Φ declines over the second interval because species arising late in that interval have less time to generate descendants during the time of elevated sampling than do species arising early in the interval. Similarly, Φ increases over the fifth interval because species arising late in the interval have a higher probability of having successors present during intervals of normal ψ . These patterns are mirrored in the actual probabilities of a sampled sister taxon arising in each interval-slices simply because λ is constant here (Fig. 7B). Finally, it is the absolute shifts in ψ that seem to be important: simply doubling low ψ has relatively little impact ψ never is high enough to greatly improve the probability of a clade member being sampled before the clade becomes extinct.

The exact probability that an earlier-sampled species or clade diverges from other sampled members of the clade in any interval-slice reflects both ψ at that point in time (i.e., the probability of sampling a predecessor) and subsequent ψ (i.e., the probability of sampling a sister-clade). The primary shifts (Fig. 7C) strongly reflect the changing probability of sampling a predecessor, particularly when ψ is high and the probability of a sampled predecessor or the divergence of a sampled sister taxon is high. However, changes in Φ in response to reduced ψ in later intervals cause the probability of a sampled divergence date to grade from one interval to the next, rather than simply be flat throughout an interval (see, e.g., Wagner 2000). As above,

this effect is muted when ψ is low, simply because absolute values are more important than relative values here.

When ψ is high, the expected branch durations ($E[d]$) increase or decrease rapidly around intervals of depressed or elevated sampling (Fig. 7D-E) and then return to “normal.” When ψ is low, shifts in $E[d]$ are extended over much longer period of time. Reduced sampling lengthens expected branch durations for taxa evolving well after the low- ψ interval, whereas elevated sampling reduces expected branch durations for taxa evolving well after the low- ψ interval. Indeed, at the lowest ψ used here, the effects of the elevated ψ interval on $E[d]$ extend up to taxa appearing during (or just before) the low- ψ interval! This stated, the effects are fairly muted at low- ψ , at least when ψ varies by only factors of two. Finally, simulations indicate that our estimates of Φ accurately predict the probability of sampling at least one descendant of any given divergence when ψ varies, regardless of the typical rate.

Effects of shifts in diversification rates (λ and μ)

I use five shifts in diversification rates to stereotype five basic macroevolutionary scenarios:

1. a burst of high speciation with constant extinction;
2. an interval of elevated continuous extinction and constant origination;
3. an interval ending with elevated pulsed extinction and constant origination;
4. an interval ending with elevated pulsed extinction followed by a rebound interval of elevated origination;

Figure 8. Probabilities of sampling clades of unknown size (Φ) under exemplar models of diversification. Origination, extinction and sampling rates are per Myr, with each interval 2.5 Myr long. Each Φ_i is for a 100-kyr slice. Black lines indicate Φ based on text equation 9 and the rates of origination (λ , in light gray) and extinction (μ , in dark gray) of that time-slice and subsequent time-slices. For each curve, one of five sampling rates (ψ) is assumed for the entire duration. For pulsed extinction, the gray bar shows the average stage rate, with $\mu_i=0.4$ for the first 24 slices and $\mu_i=10.4$ for the final interval-slice, giving average $\mu = 0.8$. Yellow dots represent the results of 10000 simulations and give the proportion of runs in which a lineage appearing at that time ultimately generates a sampled specimen given subsequent diversification.

I also include constant rates of λ and μ (Figs. 8A, 9A, 10A, 11A). This provides a comparison for assessing the importance of shifting rates and also allows comparison with both Bapst's (2013) and Didier et al.'s (2017) estimates of Φ (Table 2).

Table 2. The probability of sampling a clade of unknown size given origination and extinction = 0.4 and different levels of sampling (ψ). Φ_B gives the global estimate for a clade given Bapst (2013). Φ_D gives the global estimate given Didier et al (2017). med. Φ_i gives the median 100-kyr estimate that allows for variation in diversification and sampling.

ψ	Φ_B	Φ_D	med. Φ_i
0.400	0.6180	0.6180	0.6259
0.126	0.4260	0.4260	0.4297
0.040	0.2702	0.2701	0.2716
0.013	0.1628	0.1627	0.1618
0.004	0.0955	0.0951	0.0867

Constant Diversification and Sampling – When diversification and sampling are constant over time, the estimates of Φ that allow for temporal variation in diversification and sampling rates yield are nearly identical to estimates assuming constant rates (Table 2). The greatest difference is seen when ψ is very low. As discussed below, this reflects the fact that the edge-effect of not allowing sampling 13+ intervals later begins to affect the probability of sampling sister-taxa at very low-levels of sampling.

Intervals of Elevated Origination – Here, λ doubles in the second interval, and then decreases by 1.414 times over both of the next two intervals to return to the “normal” level. Elevating λ raises Φ not just in the intervals of high λ , but also in preceding intervals (Fig. 8B). The latter effect becomes more pronounced as ψ decreases: at $\psi=0.4$ (50% taxon-sampling given $\mu=0.4$ and $P[\text{taxon-sampling}] = \frac{\psi}{\mu+\psi}$; Foote 1996), Φ at the onset of the interval preceding diversification is close to that expected given normal rates (Fig. 8A); however, given $\psi=0.004$ (~0.99% taxon-sampling given $\mu=0.4$), Φ at the onset of the preceding interval is twice that expected given normal rates. Accordingly, the probability of a *sampled* clade diverging in the interval preceding a radiation is greater than the probability of a sampled clade diverging after that radiation even if the origination rates themselves are the same before and after (Fig. 9B).

The effects of elevated λ on expected branch durations and stratigraphic gaps within a tree ($E[d]$)

also are more dramatic when ψ is low than when ψ is high (Figs. 10B, 11B). When ψ is high, expected sampled predecessors make long branch durations and stratigraphic gaps improbable given either high or low λ (see, e.g., Foote 1996). On the other hand, when ψ is low, divergence

Figure 11. Expected branch durations ($E[d]$) for a species first appearing in an interval-slice or for an unsampled ancestor diverging in that interval-slice. $E[d]$ is the branch duration where $P[d|\lambda, \mu, \psi] = 0.5$. Note that $E[d]$ at low ψ allows for several preceding intervals with “normal” diversification and sampling rates.

486 times usually are truncated by sampled sister taxa and the increased probability of sampled sister
 487 taxa when or before λ is high reduces expected branch durations. Thus, the effect of elevated λ
 488 on $E[d]$ is more pronounced under low ψ than under high ψ .

Interval of Continuous Elevated Extinction – Here, μ is twice normal for one interval. Elevated μ causes both Φ and expected sampled sister taxa to decrease in the intervals leading up to it (Figs. 7C, 8C). This reflects the decreased probability of a lineage having numerous and/or long-lived progeny. As ψ decreases, the effect of elevated μ on Φ extends further back in time before the extinction interval. However, Φ increases as the extinction interval progresses, reflecting the fact that the increasing probability of extant species having surviving successors that might be sampled in subsequent intervals.

Intervals of elevated extinction increase the probability of stratigraphic gaps (Fig. 9C) and thus inflate expected branch durations and stratigraphic gaps for species first appearing either before or after the interval of extinction (Fig. 10C, 11C). This reflects the increased probability of sister-taxa going extinct before generating a sampled species. The effect becomes more pronounced as ψ decreases and the probability of setting a divergence time with a sampled predecessor decreases: not only is $E[d]$ higher during the extinction interval, $E[d]$ is also inflated in earlier and later intervals (Fig. 8C vs. 8A at the same ψ).

Interval Ending in Pulsed Elevated Extinction - There is considerable evidence that some major extinctions occur as pulses at the end of intervals (Marshall 1995b; Jin et al. 2000; Wang and Marshall 2004; Wang et al. 2014). Therefore, I model pulsed extinction here by setting $\mu_i=10.4$ for the final interval-slice to after $\mu_i=0.4$ for the first 24 interval-slices. The probability of extinction for an individual species present at the outset of the high-extinction interval is the same as in the continuous elevated extinction example (0.86), but with the bulk of the extinction

concentrated in the final interval-slice. This generates a similar decline in Φ and expected sampled divergences as we see given elevated continuous extinction (Figs. 10-11D vs. 10-11C), but with the decline is now offset so that the minimum values are at the end of the extinction interval rather than at the outset. Correspondingly, the rebound to “normal” Φ and expected sampled divergences is immediate rather than protracted. The differences in both offset and the rebound reflect species appearing in the middle of the interval not having appreciably better probabilities of having surviving successors than do species at the outset of the interval. The amount of time with depressed Φ prior to the extinction also is lower with pulsed extinction than with continuous extinction of the same magnitude, particularly when ψ is low. The effect on expected branch durations and stratigraphic gaps at any given sampling level ψ is muted under pulsed extinction relative to continuous extinction (Figs. 9D vs. 9C and 10D vs. 10C). Not only are the maximum deviations from “normal” lower, but the departures from normal begin and end closer to the extinction event.

Pulsed Elevated Extinction Followed by Rebound in Origination - Intervals following elevated extinction often show rebounds with elevated origination (e.g., Miller and Sepkoski 1988; Lu et al. 2006; Foote et al. 2018). The probability of sister-taxa evolving in the last interval and then surviving the extinction are the same as in the prior example. However, there now is a much higher probability that any species evolving prior to the extinction that do have surviving successors will generate a sampled descendant, which dampens the drop in Φ (Figs. 8E-9E vs. Figs. 8D-9D). This results in a mixing of the expected phylogenetic patterns preceding extinctions and radiations: the extinction reduces the probability of sampled sister taxa diverging prior to the extinction and subsequent, which offsets the pre-radiation decrease in expected

branch durations (Figs. 10E-11E vs. Figs. 10B-11E); however, the elevated probability of sampling the successors of any survivors dampens the expected increase in branch durations induced by the extinction (Figs. 10E-11E vs. Figs. 10D-11D). As in other scenarios, low ψ results in more time elapsing after the rebound before we return to “normal” branch durations and stratigraphic gaps. In particular, $E[d]$ decreases going back in time (Fig. 11E) until the rebound interval, and then rapidly increases. Going further back in time, $E[d]$ increases slightly rather than decreases (as expected when there is no rebound), with the time of increase greater at low ψ than at high ψ .

Complete Extinction (or End of Study Edge-Effect) – Whole clades might go extinct during pulsed extinctions, which ends the chance to sample subsequent successors. For example, in a phylogenetic analysis of non-avian dinosaurs or ammonites from the Cretaceous, there is no opportunity for an unsampled Maastrichtian species to leave Paleocene or Eocene species that would document that particular divergence. Therefore, divergences that are not sampled in the Maastrichtian are never sampled. This has strong effects on branch durations. Even if λ , μ and ψ are static prior to the extinction event, Φ declines markedly before the extinction (Fig. 8F). Because divergences generating sampled sister taxa become rarer (Fig. 9F), expected branch durations increase as taxon first appearances approach the extinction (Figs. 10F-11F).

The effect of ψ on Φ and expected sampled sister taxa seen for elevated extinction rates above becomes even more pronounced given total extinction. Under high ψ , Φ begins to decline notably only about 1.5 intervals (= 1.5 times the average species duration) before the extinction. However, at the lowest ψ , Φ is only half of “normal” (Fig. 8F vs. Fig. 8A) six intervals (again,

six times the average species duration) before the extinction. We see a corresponding effect on expected branch durations (Fig. 11F). When ψ is high, $E[d]$ is low until just before the extinction: even though we have little chance of sampling late divergences in any way, we have a high probability of sampling the predecessors of those late-appearing species that we do sample. When ψ is low and we do not expect to sample many direct predecessors, $E[d]$ just prior to the extinction can be nearly twice $E[d]$ under the same “normal” diversification and sampling rates (Fig. 11E vs. 11A), simply because of our inability to sample survivors of the extinction.

The sudden extinction scenario here is the same as edge-effect that excluding post-Katian species induces on the expected branch durations for an analysis of pre-Hirnantian gastropods only. As also noted above, this edge effect can be seen in Table 2. At the two lowest sampling levels, median Φ given homogeneous λ , μ and ψ is lower when we allow for temporal variation than when we do not. At the second lowest level (where we expect to sample about 3% of taxa), Φ_i drops below the “global” estimate over 18 intervals before the end of the calculations. At the lowest ψ , Φ_i is less than the estimates assuming constant diversification and sampling at least 30 intervals before the conclusion.

Discussion

Simulations illustrated in Figs. 7-8 and Supplementary Fig. S4 indicate that the approach here does well at predicting the probability of sampling a species and/or any of its descendants given “future” sampling and diversification. This reflects Eq. 6 and its building blocks adequately converting λ_i and μ_i to S_i , the expected summed durations of a lineage or clade members within an interval-slice. As this is the most complicated portion of estimating probabilities of

unsampled branch durations, this gives us some confidence that the approach will work with real data using even more complex interval-to-interval variation in diversification and sampling. Therefore, I will discuss first some implications for tree-based studies, and then outline some further modifications that we can make to accommodate evolutionary scenarios not considered above.

Shifts in Net-Diversification Rates and Shifts in Sampling Rates have Similar Effects on the Prior Probability of Sampled Sister-Taxa

Comparing Fig. 7 to Figs. 8-11 corroborates a conclusion that necessarily follows from the works of Foote et al. (1999), Bapst (2013) and Didier et al. (2017): increasing net-diversification rates has the same general effect as increasing sampling rates whereas decreasing net-diversification rates has the same general effect as decreasing sampling rates. The primary difference in how net-diversification and sampling affect branch durations is that whereas both affect the probability of sampling sister-taxa, only sampling rates affect the probability of sampling predecessors. A corollary to this is that the relative effects of shifts in diversification rates on expected branch durations and divergence times increase as overall sampling decreases: if sampled predecessors only infrequently terminate branch durations, then we require changes in λ and/or μ to increase or decrease the probability of sampling a sister taxon. Conversely, if we had 100% taxon-sampling, then sister taxa would never set divergence times as we would find all predecessors.

Phylogenetic Signor-Lipps Effect

Elevated extinction rates over short periods of time coupled with imperfect sampling create “gradual” series of last appearances in the fossil record (i.e., the Signor-Lipps effect; Signor and Lipps 1982, Raup 1986). This extends to phylogenetic patterns: because elevated extinction decreases the probability that any descendant of a divergence will ever be sampled, we expect there to be fewer nodes on phylogenies leading up to major extinctions even if rates of λ , μ and ψ are static up to that extinction. The magnitude of the effect increases both with the severity of the extinction and with the paucity of the fossil record. The latter is particularly important. Consider ammonites and non-avian dinosaurs, both of which are entirely eliminated by the K/Pg extinction. Because ammonites almost certainly have much higher ψ than do dinosaurs (Foote and Sepkoski 1999; Bapst et al. 2016), we expect the K/Pg to induce less “distortion” in the branch durations and stratigraphic gaps within ammonite phylogeny than within dinosaur phylogeny (Figs 7F, 8F, 9F).

Dinosaurs offer a possible warning case here. Sakamoto et al. (2016) note a decreasing frequency of nodes in non-avian dinosaurs phylogeny through the latter half of the Cretaceous. They attribute this to prolonged decline in λ . However, we expect at least part of this to be due to the K/Pg extinction eliminating non-avian dinosaurs and creating a phylogenetic Signor-Lipps effect that reduces Late Cretaceous Φ and thus expected nodes among Late Cretaceous taxa. The question then becomes, how far back in time might this effect have stretched? Answering this requires first answering another question: what are typical sampling rates for dinosaur? Starrfelt and Liow (2016) use the TRiPS method to estimate the overall sampling probability of dinosaur species to be greater than 0.5. As sampling probability equals $\frac{\psi}{\mu+\psi}$ (Foote 1996), this implies that

sampling rates exceed extinction rates and that any Signor-Lipps effect on phylogeny will be restricted to the latest Maastrichtian (Figs. 8F-11F). However, Bapst et al. (2016; see also Benson et al. 2018) use Foote's (1997) method to estimate that dinosaur ψ as about 60 times lower than dinosaur μ . This is most comparable to the expectations given the lowest ψ used here. Bapst et al. estimated $\mu=0.935$, which in turn implies an average dinosaur species duration of 1.1-myr. At the lowest sampling levels examined above, the phylogenetic Signor-Lipps effect extends back 30 times the average species lifetime. For dinosaurs, that would mean reduced sampling of sister-taxa extending back over 30-myr, i.e., to the Campanian. Relative sampling could be even lower than that for dinosaurs. Distributions of ψ among contemporaneous mammal species follow lognormal distributions (Wagner and Marcot 2013; see also Foote 2016) and the sampling rates estimated by methods such as Foote's 1997 one typically are 10 to 100 times higher than median (=geometric mean) lognormal rates. If so, then the Signor-Lipps effect on Φ caused by the K/Pg extinction could be more severe than any case illustrated here. On the other hand, interest in extinction patterns leading up to the K/Pg might create unusually high ψ in the Maastrichtian, which in turn should increase the proportions of nodes found in the Late Cretaceous (see, e.g., Starrfelt and Liow 2016: Fig. 1). This would be similar to a "Pull of the Recent" (Raup 1979) effect where the high sampling of extant taxa increases the representation of recent divergences on phylogenies (Heath et al. 2014). Thus, assessing the effects of the total extinction of non-avian dinosaurs requires not just the typical sampling rates, but also the heterogeneity in those sampling rates over time.

Low Sampling will “Blur” Radiations back in Time

When using appearance or occurrence data, incomplete sampling has the same effect on the apparent timing of radiations as it does on extinctions, albeit in reverse, by making events look more gradual than they truly were (i.e., the Jaanusson Effect or Sppil-Rongis Effect; Jaanusson 1976; Marshall 1995a, 1998). However, sampling induces the *opposite* effect on the expected frequencies of sampled divergences (= nodes on a cladogram) during radiations, as we expect the frequency of nodes to increase before λ increases (Figs. 8B, 9B). As a corollary, the expected branch durations/stratigraphic gaps ($E[d]$; Figs. 10B, 11B) should decrease before λ increases.

The blurring effect becomes more pronounced as ψ decreases. To envision why this would be, suppose that we are examining two Ordovician echinoderm and brachiopod clades that show elevated net-diversification during the Darriwilian. The Jaanusson / Sppil-Rongis effect should smear the first appearances of taxa appearing during the radiation towards later Ordovician stages (e.g., the Sandbian or even Katian). This should be more true for echinoderms than for brachiopods because per-taxon-sampling rates generally are higher for brachiopods than for echinoderms (Foote and Sepkoski 1999). However, unsampled Dapingian ancestral species will have higher probabilities of generating sampled descendants than will unsampled Floian ancestral species: Dapingian species have greater probabilities than do Floian species of having successors that radiate in the Darriwilian. The effect also should be more dramatic for echinoderms than for brachiopods. Compared to echinoderms, we should sample more Dapingian brachiopod species (or near-relatives of those species) that give rise descendants during the Darriwilian. In other words, we should be sampling more of both the “dark gray/dark blue” and “light gray/yellow” lineages from Fig. 1 for Dapingian brachiopods than for Dapingian

echinoderms. As a corollary, compared to brachiopods, echinoderms should have more Dapingian nodes that link two Darriwilian nodes rather than one or more sampled taxa. This results in reduced branch durations and individual stratigraphic gaps *per branch* while at the same time permitting the Jaanusson / Sppil-Rongis effect with sampled taxa.

An additional corollary to that elevated origination rates following elevated extinction rates will partially offset the effects of extinction on expected branch durations and nodes within a tree (Figs. 8E-11E). Thus, simple summaries of how branch durations and/or nodes from phylogenies are distributed over time might miss major turnovers. Instead, we need to compare the distributions of branch durations/stratigraphic gaps: relative to other intervals, species and branches dated to the rebound interval should show a greater proportion of both relatively short and relatively long branches compared to “normal” intervals with the same expected branch durations. Basically, we should either have a short branch duration confined to the rebound interval or a much longer one stretching back well into the extinction interval or even earlier. Alternatively, distributions of these parameters from particular portions of phylogenies (e.g., Soul and Friedman 2017) or simply “traditional” quantitative methods such as paleobiologists have been developing for decades will be of more use here.

Implications for Morphological Rate Studies and Tip-Dating

It is common for morphological disparity to peak early in clade histories (Hughes et al. 2013). Tip-dating approaches should be prone to implying deep divergence times for such clades, in order to account for the large numbers of differences among early taxa. However, if those early bursts of disparity accompany elevated rates of diversification, then deep divergence times will

be improbable given birth-death-sampling models. This could add considerable power to tree-based studies of shifting rates of morphological change, and offer an improvement on prior assessments of the “deep unsampled divergences” alternative than sampling alone (e.g., Wagner 1995b; Ruta et al. 2006; Hopkins and Smith 2015). Moreover, appropriate priors on branch durations should at least somewhat mitigate the effects of correlated character change, which likely is common among fossil taxa (Wagner and Estabrook 2015), and which should mislead tip-dating methods by implying numerous character changes when only a single change occurred (see, e.g., Pagel and Meade 2006).

Implications for Analyses of Stratigraphic Gaps within Phylogenies

As noted above, the branch durations implicit to phylogenies correspond to gaps in stratigraphic sampling within clades. Numerous studies contrast the stratigraphic gaps implicit to inferred phylogenies from different taxa or intervals of time (e.g., Benton and Storrs 1994; Hitchin and Benton 1997; Wills 1999, 2007; Benton et al. 2000; O'Connor et al. 2011). O'Connor et al. (2016) assess a variety of reasons why differences might exist other than sampling, although they did not include differences in diversification rates. Bapst (2013) shows that differences in diversification offer another reason why two clades with otherwise similar preservation potential will have phylogenies with different summed branch durations and thus stratigraphic gaps. To this, we need to add heterogeneity in diversification parameters as a reason why two clades with similar preservation potential *and* similar general diversification dynamics should have different summed stratigraphic gaps.

Accommodating Pulsed Turnover

The estimates of Φ and expected branch duration probabilities presented above assume that origination and extinction are always happening. Pulsed turnover models posit that origination and/or extinction might be concentrated at the ends/onsets of what we now recognize as stratigraphic boundaries (Foote 2005). Purely pulsed extinction with continuous origination posits pure-birth for diversification within an interval. Now the probability of one species having $K-1$ descendants after duration t is:

$$P[K | \lambda, t] = e^{-\lambda t} \times (1 - e^{-\lambda t})^{K-1} \quad (\text{Eq. 12})$$

(Yule 1925). Equations 2-5 still give the probabilities of 0...N species surviving given some extinction rate. Note that Φ will be much higher at any given combination of λ , μ and ψ given pulsed extinction than given continuous extinction because $P[0 \text{ species}] = 0$ until the very end of the interval. Thus, an interval of length T has a minimum $L_{\text{my}}=T$ rather than nearly zero. For pulsed origination, pure-birth can be used for diversification at the outset of the interval. If both birth and death are pulsed, then Φ is maximized because all lineages will persist for (essentially) the entire interval. This is essentially a corollary of Foote's (2005) demonstration that pulsed origination and extinction increases the proportion of taxa that we should first/last observe in the first/last intervals at any given ψ . Of course, pulsed origination also hugely simplifies searching for divergence times: all branching would be concentrated in a small number of short intervals under a pure turnover-pulse model.

Finally, we might have mixed models in which there is some background rate and then some short-term pulsed rate. This is illustrated above under the pulsed extinction models (Figs. 8D-E –

11D-E). To some extent using time-slices finer than intervals as done here obviates the entire issue: the pure pulsed models then would have only the first and/or last time slices with rates above zero.

Accommodating within-interval variation in diversification and sampling among clade members We can accommodate rate variation among taxa within time intervals by using hyperpriors on the rate distributions (see Heath 2012). For example, sampling distributions among contemporaneous taxa often fit lognormal distributions (Wagner and Marcot 2013; Foote 2016). One could use the rates from midpoints of the four quartiles and calculate the average Φ , the average probability of a sampled divergence and the average probability of 0 finds given those four rates (see, e.g., Yang 1994). It is less clear what the appropriate distributions should be for λ and μ . Although paleontologists have modelled variation in extinction among intervals (e.g., Raup 1991; Wang 2003), we have not yet modelled variation in diversification so among taxa within intervals. Heath et al. (2017) use exponential distributions for diversification hyperpriors. However, gamma distributions might be more appropriate than exponentials, as continuous origination and extinction rates are Poisson processes and the gamma distribution is the conjugate prior for Poisson and related processes. Pulsed extinction is consistent with a binomial process (i.e., some proportion of species survive), and the conjugate prior there is a beta distribution. Yet another possibility is that λ and μ reflect occupancy patterns within species and thus either follow or are affected by distributions describing occupancy such as lognormals (Foote 2016). This latter possibility also means that λ and μ should co-vary among closely related species (see, e.g., Roy et al. 2009). If so, then skyline models allowing for heritable shifts in diversification rates might be most appropriate (Stadler et al. 2013; see also Stadler et al.

2018). Regardless, a very important question for paleobiologists to answer in the near future is: exactly what distributions do best describe among-taxon variation in λ and μ ?

Other issues

Stadler et al. (2018) note that anagenetic transformation of predecessors into derived species, either during bifurcating cladogenesis (Fig. 1C) or within lineages, will estimates of origination and extinction implied by the stratigraphic ranges of named species. This in turn will alter our ideas on the probability of sampled sister taxa diverging. Many paleobiological studies suggest that anagenetic transformation on phylogenies is uncommon, but not non-existent (Wagner and Erwin 1995; Polly 1997; Pardo et al. 2008; Bapst and Hopkins 2017). Stadler et al. (2018) suggest using two parameters, λ_B and λ_A , for branching and anagenetic speciation, respectively. One then could then modify either parameter λ_A over the course of Markov Chain Monte Carlo searches, constraining $\lambda_{B \cdot i} + \lambda_{A \cdot i} = \lambda_i$. Correspondingly, we would need to separate “true” extinction (μ_T) from pseudoextinction rate (also $\lambda_{A \cdot i}$) so that $\mu_{T \cdot i} + \lambda_{A \cdot i} = \mu_i$. Under this approach, if the trees maximizing the likelihood of morphological change models also imply copious anagenetic change, then those trees will not have reduced priors due to not having the sampled sister-taxa predicted by λ (or $\lambda_{1 \dots N}$ for N intervals) as they instead will use the lower λ_B and higher λ_A .

Conclusion

As Bapst (2013) stresses, the probabilities of branch durations and divergence times represent a three-rate problem, with diversification and sampling affecting expected branch durations and stratigraphic gaps within phylogenies of fossil taxa. Heterogeneity in diversification over time

also affects these expectations. The method presented herein allows paleobiologists to accommodate temporal variation in diversification and sampling that so many paleobiological studies document, and potentially overcomes a chief criticism of birth-death-sampling models (e.g., Marshall 2017). Given that current methods for modeling diversification and sampling with fossil data are almost certainly more advanced than are our current methods for modeling anatomical character evolution, the approach developed here provides paleobiologist with a much-needed independent test of divergence times implicit to alternate phylogenetic hypotheses, and thus an additional control when testing ideas about relationships or patterns of evolution across phylogeny.

Acknowledgements

I thank R. Warnock for copious discussion about birth-death-sampling models in all of their different guises. D. Bapst, M. Foote, W. Kiessling, M. Sakamoto and an anonymous reviewer provided very helpful comments. For Katian gastropod data entered into the Paleobiology Database, I thank S. Holland and B. Kröger. This is Paleobiology Database Publication #_____

References

- Alroy, J. 2014. Accurate and precise estimates of origination and extinction rates. *Paleobiology* 374-397.
- Bapst, D. W. 2013. A stochastic rate-calibrated method for time-scaling phylogenies of fossil taxa. *Methods in Ecology and Evolution* 4:724-733.
- . 2016. paleotree. program distributed by the <https://github.com/dwbapst/paleotree>.

- 805 Bapst, D. W., and M. J. Hopkins. 2017. Comparing cal3 and other a posteriori time-scaling
806 approaches in a case study with the pterocephaliid trilobites. *Paleobiology* 43:49-67.
- 807 Bapst, D. W., A. M. Wright, N. J. Matzke, and G. T. Lloyd. 2016. Topology, divergence dates,
808 and macroevolutionary inferences vary between different tip-dating approaches applied to
809 fossil theropods (Dinosauria). *Biology Letters* 12:20160237.
- 810 Benson, R. B. J., G. Hunt, M. T. Carrano, and N. Campione. 2018. Cope's rule and the adaptive
811 landscape of dinosaur body size evolution. *Palaeontology* 61:13-48.
- 812 Benton, M. J., and G. W. Storrs. 1994. Testing the quality of the fossil record: paleontological
813 knowledge is improving. *Geology* 22:111 - 114.
- 814 Benton, M. J., M. A. Wills, and R. Hitchin. 2000. Quality of the fossil record through time.
815 *Nature* 403:534 - 537.
- 816 Bergström, S. M., X. Chen, J. C. Gutiérrez-Marco, and A. Dronov. 2009. The new
817 chronostratigraphic classification of the Ordovician System and its relations to major
818 regional series and stages and to $\delta^{13}\text{C}$ chemostratigraphy. *Lethaia* 42:97-107.
- 819 Connolly, S. R., and A. I. Miller. 2001. Joint estimation of sampling and turnover rates from
820 fossil databases: capture-mark-recapture methods revisited. *Paleobiology* 27:751 – 767.
- 821 Didier, G., M. Fau, and M. Laurin. 2017. Likelihood of tree topologies with fossils and
822 diversification rate estimation. *Systematic Biology* 66:964-987.
- 823 Finarelli, J. A., and L. H. Liow. 2016. Diversification histories for North American and
824 Eurasian carnivorans. *Biological Journal of the Linnean Society* 118:26 – 38.
- 825 Fisher, D. C. 1994. Stratocladistics: morphological and temporal patterns and their relation to
826 phylogenetic process. Pp. 133 - 171 in L. Grande and O. Rieppel, eds. *Interpreting the*

- 827 hierarchy of nature - from systematic patterns to evolutionary process theories. Academic
828 Press, Orlando, Florida.
- 829 Foote, M. 1996. On the probability of ancestors in the fossil record. *Paleobiology* 22:141 - 151.
830 —————. 1997. Estimating taxonomic durations and preservation probability.
831 *Paleobiology* 23:278 - 300.
- 832 —————. 2001. Inferring temporal patterns of preservation, origination, and
833 extinction from taxonomic survivorship analysis. *Paleobiology* 27:602 - 630.
- 834 —————. 2003. Origination and extinction through the Phanerozoic: a new
835 approach. *The Journal of Geology* 111:125 - 148.
- 836 —————. 2005. Pulsed origination and extinction in the marine realm. *Paleobiology*
837 31:6-20.
- 838 —————. 2016. On the measurement of occupancy in ecology and paleontology.
839 *Paleobiology* 42:707 - 729.
- 840 Foote, M., R. A. Cooper, J. S. Crampton, and P. M. Sadler. 2018. Diversity-dependent
841 evolutionary rates in early Palaeozoic zooplankton. *Proceedings of the Royal Society B:*
842 *Biological Sciences* 285:20180122.
- 843 Foote, M., J. P. Hunter, C. M. Janis, and J. J. Sepkoski, Jr. 1999. Evolutionary and
844 preservational constraints on origins of biologic groups: divergence times of eutherian
845 mammals. *Science* 283:1310 - 1314.
- 846 Foote, M., and J. J. Sepkoski, Jr. 1999. Absolute measures of the completeness of the fossil
847 record. *Nature* 398:415 - 417.

- 848 Friedman, M. 2010. Explosive morphological diversification of spiny-finned teleost fishes in
849 the aftermath of the end-Cretaceous extinction. *Proceedings of the Royal Society B:*
850 *Biological Sciences* 277:1675-1683.
- 851 Gavryushkina, A., T. A. Heath, D. T. Ksepka, T. Stadler, D. Welch, and A. J. Drummond. 2016.
852 Bayesian Total-Evidence Dating reveals the Recent crown radiation of penguins. *Systematic*
853 *Biology* 66:57-73.
- 854 Gradstein, F. M., J. G. Ogg, and M. Schmitz. 2012. *The geologic time scale 2012*. Elsevier,
855 Amsterdam.
- 856 Halliday, T. J. D., and A. Goswami. 2016. The impact of phylogenetic dating method on
857 interpreting trait evolution: a case study of Cretaceous–Palaeogene eutherian body-size
858 evolution. *Biology Letters* 12:20160051.
- 859 Heath, T. A. 2012. A Hierarchical Bayesian model for calibrating estimates of species
860 divergence times. *Systematic Biology* 61:793-809.
- 861 Heath, T. A., J. P. Huelsenbeck, and T. Stadler. 2014. The fossilized birth–death process for
862 coherent calibration of divergence-time estimates. *Proceedings of the National Academy of*
863 *Sciences* 111:E2957–E2966.
- 864 Heath, T. A., A. M. Wright, and W. Pett. 2017. Total-Evidence Analysis of Fossil and Extant
865 Taxa under the Fossilized Birth-Death Process. Pp. 1 - 31 in S. Höhna, M. J. Landis and T.
866 A. Heath, eds. *Phylogenetic Inference Using RevBayes*. *Current Protocols in Bioinformatics*,
- 867 Hitchin, R., and M. J. Benton. 1997. Congruence between parsimony and stratigraphy:
868 comparisons of tree indices. *Paleobiology* 23:20 - 32.

- 869 Hopkins, M. J., and A. B. Smith. 2015. Dynamic evolutionary change in post-Paleozoic
870 echinoids and the importance of scale when interpreting changes in rates of evolution.
871 *Proceedings of the National Academy of Sciences* 112:3758–3763.
- 872 Huelsenbeck, J. P., and B. Rannala. 1997. Maximum likelihood estimation of topology and
873 node times using stratigraphic data. *Paleobiology* 23:174 - 180.
- 874 Hughes, M., S. Gerber, and M. A. Wills. 2013. Clades reach highest morphological disparity
875 early in their evolution. *Proceedings of the National Academy of Sciences* 110:13875-
876 13879.
- 877 Jaanusson, V. 1976. Faunal dynamics in the middle Ordovician (Viruan) of Balto-Scandia. Pp.
878 301-326 in M. G. Bassett, ed. *The Ordovician System: Proceedings of a palaeontological*
879 *association symposium*. University of Wales Press and National Museum of Wales, Cardiff.
- 880 Jin, Y. G., Y. Wang, W. Wang, Q. H. Shang, C. Q. Cao, and D. H. Erwin. 2000. Pattern of
881 marine mass extinction near the Permian-Triassic boundary in South China. *Science* 289:432
882 - 436.
- 883 Lewis, P. O. 2001. A likelihood approach to estimating phylogeny from discrete morphological
884 character data. *Systematic Biology* 50:913 - 925.
- 885 Liow, L. H., and J. A. Finarelli. 2014. A dynamic global equilibrium in carnivoran
886 diversification over 20 million years. *Proceedings of the Royal Society B: Biological*
887 *Sciences* 281:20132312.
- 888 Liow, L. H., and J. D. Nichols. 2010. Estimating rates and probabilities of origination and
889 extinction using taxonomic occurrence data: capture-mark-recapture (CMR) approaches. Pp.
890 81 - 94 in J. Alroy and G. Hunt, eds. *Quantitative methods in paleobiology*. *Paleontological*
891 *Society*, New Haven, Connecticut.

- 892 Lu, P. J., M. Yogo, and C. R. Marshall. 2006. Phanerozoic marine biodiversity dynamics in
893 light of the incompleteness of the fossil record. *Proceedings of the National Academy of*
894 *Sciences of the United States of America* 103:2736-2739.
- 895 Marshall, C. R. 1995a. Stratigraphy, the true order of species' originations and extinctions, and
896 testing ancestor-descendant hypotheses among Caribbean bryozoans. Pp. 208 - 236 in D. H.
897 Erwin and R. L. Anstey, eds. *New approaches to studying speciation in the fossil record*.
898 Columbia University Press, New York.
- 899 —————. 1995b. Distinguishing between sudden and gradual extinctions in the
900 fossil record: predicting the position of the iridium anomaly using the ammonite fossil record
901 on Seymour Island, Antarctica. *Geology* 23:731 - 734.
- 902 —————. 1998. Determining stratigraphic ranges. Pp. 23-53 in C. R. C. Paul and S.
903 K. Donovan, eds. *The adequacy of the fossil record*. Wiley, New York.
- 904 —————. 2017. Five palaeobiological laws needed to understand the evolution of
905 the living biota. *Nature Ecology & Evolution* 1:0165.
- 906 Matzke, N. J., and R. B. Irmis. 2018. Including autapomorphies is important for paleontological
907 tip-dating with clocklike data, but not with non-clock data. *PeerJ* 6:e4553.
- 908 Matzke, N. J., and A. M. Wright. 2016. Inferring node dates from tip dates in fossil Canidae:
909 the importance of tree priors. *Biology Letters* 12:2016032.
- 910 Miller, A. I., and J. J. Sepkoski, Jr. 1988. Modeling bivalve diversification: the effect of
911 interaction on a macroevolutionary system. *Paleobiology* 14:364 - 369.
- 912 O'Connor, A., C. Moncrieff, and M. A. Wills. 2011. Variation in stratigraphic congruence
913 (GER) through the Phanerozoic and across higher taxa is partially determined by sources of
914 bias. *Geological Society, London, Special Publications* 358:31-52.

- 915 O'Connor, A., and M. A. Wills. 2016. Measuring stratigraphic congruence across trees, higher
916 taxa, and time. *Systematic Biology* 65:792-811.
- 917 Pagel, M. D., and A. Meade. 2006. Bayesian analysis of correlated evolution of discrete
918 characters by Reversible-Jump Markov Chain Monte Carlo. *The American Naturalist*
919 167:808-825.
- 920 Pardo, J. D., A. K. Huttenlocker, and J. D. Marcot. 2008. Stratocladistics and evaluation of
921 evolutionary modes in the fossil record: an example from the ammonite genus
922 *Semiformiceras*. *Palaeontology* 51:767-773.
- 923 Peters, S. E., and M. Foote. 2001. Biodiversity in the Phanerozoic: a reinterpretation.
924 *Paleobiology* 27:583 – 601.
- 925 Polly, P. D. 1997. Ancestry and species definition in paleontology: a stratocladistic analysis of
926 Paleocene-Eocene Viverravidae (Mammali, Carnivora) from Wyoming. *Contributions from*
927 *the Museum of Paleontology, the University of Michigan* 30:1 - 53.
- 928 Raup, D. M. 1972. Taxonomic diversity during the Phanerozoic. *Science* 177:1065 - 1071.
929 ————. 1979. Biases in the fossil record of species and genera. *Bulletin of the*
930 *Carnegie Museum of Natural History* 13:85 - 91.
- 931 ————. 1985. Mathematical models of cladogenesis. *Paleobiology* 11:42 - 52.
932 ————. 1986. Biological extinction in earth history. *Science* 231:1528 - 1533.
933 ————. 1991. A kill curve for Phanerozoic marine species. *Paleobiology* 17:37 -
934 48.
- 935 Ronquist, F., S. Klopfstein, L. Vilhelmsen, S. Schulmeister, D. L. Murray, and A. P. Rasnitsyn.
936 2012. A Total-Evidence approach to dating with fossils, applied to the early radiation of the
937 Hymenoptera. *Systematic Biology* 61:973-999.

- 938 Roy, K., G. Hunt, and D. Jablonski. 2009. Phylogenetic conservatism of extinctions in marine
939 bivalves. *Science* 325:733-737.
- 940 Ruta, M., P. J. Wagner, and M. I. Coates. 2006. Evolutionary patterns in early tetrapods. I.
941 Rapid initial diversification by decrease in rates of character change. *Proceedings of the*
942 *Royal Society of London, Series B. Biological Sciences* 273:2107 - 2111.
- 943 Sakamoto, M., M. J. Benton, and C. Venditti. 2016. Dinosaurs in decline tens of millions of
944 years before their final extinction. *Proceedings of the National Academy of Sciences*
945 113:5036-5040.
- 946 Sansom, R. S., and M. A. Wills. 2017. Differences between hard and soft phylogenetic data.
947 *Proceedings of the Royal Society B: Biological Sciences* 284:20172150.
- 948 Sheehan, P. M. 1977. Species diversity in the Phanerozoic: a reflection of labor by
949 systematists? *Paleobiology* 3:325-328.
- 950 Signor, P. W., and J. H. Lipps. 1982. Sampling bias, gradual extinction patterns and
951 catastrophes in the fossil record. *Geological Society of America Special Paper* 190:291 -
952 296.
- 953 Smith, A. B. 1988. Patterns of diversification and extinction in early Palaeozoic echinoderms.
954 *Palaeontology* 31:799 - 828.
- 955 ————. 2001a. Probing the cassiduloid origins of clypeasteroid echinoids using
956 stratigraphically restricted parsimony analysis. *Paleobiology* 27:392 - 404.
- 957 ————. 2001b. Large-scale heterogeneity of the fossil record: implications for
958 Phanerozoic biodiversity studies. *Philosophical Transactions of the Royal Society of London*
959 *Series B* 356:351 - 367.

- 960 Soul, L. C., and M. Friedman. 2017. Bias in phylogenetic measurements of extinction and a
961 case study of end-Permian tetrapods. *Palaeontology* 60:169-185.
- 962 Stadler, T. 2010. Sampling-through-time in birth–death trees. *Journal of Theoretical Biology*
963 267:396-404.
- 964 Stadler, T., A. Gavryushkina, R. C. M. Warnock, A. J. Drummond, and T. A. Heath. 2018. The
965 fossilized birth-death model for the analysis of stratigraphic range data under different
966 speciation concepts. *Journal of Theoretical Biology* 447:41 - 45.
- 967 Stadler, T., D. Kühnert, S. Bonhoeffer, and A. J. Drummond. 2013. Birth–death skyline plot
968 reveals temporal changes of epidemic spread in HIV and hepatitis C virus (HCV).
969 *Proceedings of the National Academy of Sciences* 110:228-233.
- 970 Starrfelt, J., and L. H. Liow. 2016. How many dinosaur species were there? Fossil bias and true
971 richness estimated using a Poisson sampling model. *Philosophical Transactions of the Royal*
972 *Society B: Biological Sciences* 371:20150219.
- 973 Wagner, P. J. 1995a. Stratigraphic tests of cladistic hypotheses. *Paleobiology* 21:153 - 178.
- 974 —————. 1995b. Testing evolutionary constraint hypotheses with early Paleozoic
975 gastropods. *Paleobiology* 21:248 - 272.
- 976 —————. 2000a. Exhaustion of cladistic character states among fossil taxa.
977 *Evolution* 54:365 - 386.
- 978 —————. 2000b. Likelihood tests of hypothesized durations: determining and
979 accommodating biasing factors. *Paleobiology* 26:431 - 449.
- 980 Wagner, P. J., and D. H. Erwin. 1995. Phylogenetic patterns as tests of speciation models. Pp.
981 87 - 122 in D. H. Erwin and R. L. Anstey, eds. *New approaches to studying speciation in the*
982 *fossil record*. Columbia University Press, New York.

- 983 Wagner, P. J., and G. F. Estabrook. 2015. The implications of stratigraphic compatibility for
984 character integration among fossil taxa. *Systematic Biology* 64:838 - 852.
- 985 Wagner, P. J., and J. D. Marcot. 2013. Modelling distributions of fossil sampling rates over
986 time, space and taxa: assessment and implications for macroevolutionary studies. *Methods in*
987 *Ecology and Evolution* 4:703 - 713.
- 988 Wang, S. C. 2003. On the continuity of background and mass extinction. *Paleobiology* 29:455-
989 467.
- 990 Wang, S. C., and C. R. Marshall. 2004. Improved confidence intervals for estimating the
991 position of a mass extinction boundary. *Paleobiology* 30:5 - 18.
- 992 Wang, Y., P. M. Sadler, S.-z. Shen, D. H. Erwin, Y.-c. Zhang, X.-d. Wang, W. Wang, J. L.
993 Crowley, and C. M. Henderson. 2014. Quantifying the process and abruptness of the end-
994 Permian mass extinction. *Paleobiology* 40:113-129.
- 995 Wills, M. A. 1999. The congruence between phylogeny and stratigraphy: randomization tests
996 and the Gap Excess Ratio. *Systematic Biology* 48:559 - 580.
- 997 —————. 2007. Fossil ghost ranges are most common in some of the oldest and
998 some of the youngest strata. *Proceedings of the Royal Society of London, Series B.*
999 *Biological Sciences* 274:2421 - 2427.
- 1000 Wright, A. M., G. T. Lloyd, and D. M. Hillis. 2016. Modeling character change heterogeneity
1001 in phylogenetic analyses of morphology through the use of priors. *Systematic Biology*
1002 65:602 – 611.
- 1003 Wright, D. F. 2017. Bayesian estimation of fossil phylogenies and the evolution of early to
1004 middle Paleozoic crinoids (Echinodermata). *Journal of Paleontology* 91:799-814.

- 1005 Wright, D. F., and U. Toom. 2017. New crinoids from the Baltic region (Estonia): fossil tip-
1006 dating phylogenetics constrains the origin and Ordovician–Silurian diversification of the
1007 Flexibilia (Echinodermata). *Palaeontology* 60:893–910.
- 1008 Yacobucci, M. M. 1999. Plasticity of developmental timing as the underlying cause of high
1009 speciation rates in ammonoids. An example from the Cenomanian Western Interior Seaway
1010 of North America. Pp. 59 - 77 in F. Olóriz and F. J. Rodriguez-Tovar, eds. *Advancing*
1011 *research on living and fossil cephalopods*. Kluwer Academic / Plenum, New York.
- 1012 Yang, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with
1013 variable rates over sites: approximate methods. *Journal of Molecular Evolution* 39:306 -
1014 314.
- 1015 Yule, G. U. 1925. A mathematical theory of evolution, based on the conclusions of Dr. J. C.
1016 Willis, F.R.S. *Philosophical Transactions of the Royal Society of London. Series B,*
1017 *Biological Sciences* 213:21 - 87.
- 1018 Zhang, C., T. Stadler, S. Klopstein, T. A. Heath, and F. Ronquist. 2016. Total-Evidence dating
1019 under the Fossilized Birth-Death Process. *Systematic Biology* 65:228-249.
- 1020

Figure 1. Hypothetical phylogeny for sampled lineages (black/light blue), unsampled predecessors (dark gray/dark blue) and unsampled sister-taxa (light gray/yellow). A) Two sampled species diverged from unsampled Species D at 310-Ma, generating branch durations (and chronostratigraphic gaps) of 6-myr for Species A and 4-myr for Species I. Increased sampling will provide earlier divergence times and shorter branch durations/chronostratigraphic gaps in one of two ways. We might sample predecessors of A or I (dark gray/dark blue). The probability of doing so is the probability of sampling an individual lineage at any one point in time. Alternatively, we might sample sister-taxa of A or I (light gray/yellow). The probability of doing so reflects origination (affecting the number of possible sister-taxa and the richness of sister-clades), extinction (affecting the richness of sister-clades and the probability of sampling individual lineages), and sampling (affecting the probability of sampling individual lineages). B) The phylogeny re-rendered as budding speciation. Each letter represents a distinct morphospecies that we would recognize prior to phylogenetic analysis. Note that the predecessors of A and I now are only part of the durations of ancestors such as Species B and Species H. However, the sum of predecessors (dark gray/dark blue) and the sum of possible “sisters” (light gray/yellow) remains unchanged. C) The phylogeny re-rendered as bifurcation speciation, with ancestral morphospecies being at least slightly altered at divergence so that Species B and Species B’ have at least one character distinguishing them. Now predecessors and ancestors nearly synonymous, save for the earliest unsampled members of Species A and I. However, the sum of predecessors and the sum of possible sister remains unchanged.

Figure 2. The probability of there being $K=0\dots 4$ successors of a species extant at time $t=0$ over an interval with duration $T=2.5$ given different combinations of origination (λ) and extinction (μ) rates. Note that rates increase by a factor of 1.414 ($2^{0.5}$) between adjacent frames. Note that the successors might include the original species at any point in time.

Figure 3. The expected time with $K=0\dots 4$ successors (including the original species) given interval duration $T=2.5$. These are the areas each curve for the corresponding K , origination (λ) and extinction (μ) in the corresponding panels in Figure 2. The sum of expected lineage durations, S , is the sum of $K \times E[T_K]$. Thus at $\lambda=0.4$ and $\mu=0.4$ (A), $E[S] = (1 \times 1.25) + (2 \times 0.313) + (3 \times 0.104) + (4 \times 0.039) \dots = 2.5$ lineage million years.

Figure 4. The effect of interval-slice length on overestimate of Φ' . Horizontal bars represent expected given 5000 simulations at the appropriate origination (λ) and extinction (μ) rates. Intervals durations are set to $T=2.5$, so at $m=250$ slices, each $t_i=0.001$. See also Supplementary Fig. S3.

Figure 5. The probability of a species extant at the outset of an interval with particular origination (λ) and extinction (μ) having $K=0\dots 4$ successors extant 0.25-myr later. The probability of the species or any successor generating a sampled species in the subsequent 0.25-myr is conditioned on the probability of $K=0\dots \infty$ survivors going into the next interval-slice.

Figure 6. Breaking down estimates of branch duration probabilities given empirically estimated rates of origination (λ), extinction (μ) and sampling (ψ). Gray bars give rates for each interval (2nd y-axis). A) The components of Φ_i , the probability of sampling a species appearing in interval-slice i or any successors of that species (i.e., a clade of unknown size). v_i (yellow dots) gives the probability of sampling that species or any successors in interval-slice i . ζ_i (light blue dots) gives the probability of sampling any successors of that species (including the original species) in interval-slice $i+1$ or anytime afterwards before the end of the Katian. Φ_i reflects the probability of doing either and therefore is $\Phi_i = 1 - ([1 - v_i] \times [1 - \zeta_i])$. B) The exact probability of a branch duration leading to a sampled species or to a node linking sampled species starting in interval-slice i . The probability of sampling a predecessor (purple) is the probability of sampling the one predecessor of a species or node extant in interval-slice i . The probability of sampling a sister-taxon (dark orange) is the probability of a divergence giving rise to a sister taxon with 1+ sampled species in interval-slice i . The probability of one or the other happening (sienna) is the probability of a phylogenetic divergence in interval-slice i given that we have some sampled species with a first occurrence after interval-slice i (or a node linking sampled species that diverges after interval-slice i). C) The probabilities of branch durations (sienna) preceding *Salpingostoma richmondensis*, which is first known from early in the *Aphelognathus divergens* conodont zone. This is broken down into the probability of duration d with no divergences to sampled sister taxa (orange) and the probability of no sampled precursors (purple). The expected branch duration (where $P[d] = 0.50$) is 3.2-myr.

Figure 7. Effects of variable sampling over time with constant origination and extinction ($\lambda=\mu=0.40$). Five median sampling rates (ψ) are illustrated, with ψ relative to the median given on by the gray bars (2nd y-axis). The six intervals of 2.5-myr are preceded by six others with median ψ . A) Φ , the probability of sampling a species or any successors given that the species is extant in an interval-slice. Yellow dots give expected Φ given 10000 simulations using ψ (and λ and μ) of the intervals of a species lifetime. B) The probability of a sampled sister-species diverging in each interval-slice. C) The probability of a phylogenetic divergence in each interval-slice from either sampling a predecessor or from the divergence of a subsequently sampled sister-taxon. D) Expected durations (d) for a species first appearing in each interval-slice. This also applies to a node that diverged in the same interval-slice. $E[d]$ is the branch duration where $P[\text{no ancestors or divergences}]$ reaches 0.50 (see Fig. 6C).

Figure 8. Probabilities of sampling clades of unknown size (Φ) under exemplar models of diversification. Origination, extinction and sampling rates are per Myr, with each interval 2.5 Myr long. Each Φ_i is for a 100-kyr slice. Black lines indicate Φ based on text equation 9 and the rates of origination (λ , in light gray) and extinction (μ , in dark gray) of that time-slice and subsequent time-slices. For each curve, one of five sampling rates (ψ) is assumed for the entire duration. For pulsed extinction, the gray bar shows the average stage rate, with $\mu_i=0.4$ for the first 24 slices and $\mu_i=10.4$ for the final interval-slice, giving average $\mu = 0.8$. Yellow dots represent the results of 10000 simulations and give the proportion of runs in which a lineage appearing at that time ultimately generates a sampled specimen given subsequent diversification.

1108 **Figure 9.** Probabilities of a divergence that ultimately generates a sampled species given the
1109 subsequent origination, extinction and sampling rates.

1110

1111 **Figure 10.** Probabilities of setting a phylogenetic divergence in a particular 100-kyr
1112 interval-slice, either by sampling a predecessor or by the divergence of a sister-taxon that is
1113 sampled in that interval-slice or in some subsequent one. The compliments of these probabilities
1114 are the probabilities of a stratigraphic gap within a phylogeny over the particular interval-slices.
1115 The product of these probabilities over any set of interval-slices gives the prior probability of a
1116 branch duration spanning that set of interval-slices.

1117

1118 **Figure 11.** Expected branch durations ($E[d]$) for a species first appearing in an interval-slice or
1119 for an unsampled ancestor diverging in that interval-slice. $E[d]$ is the branch duration where
1120 $P[d|\lambda, \mu, \psi] = 0.5$. Note that $E[d]$ at low ψ allows for several preceding intervals with “normal”
1121 diversification and sampling rates.

1122

1123

Table 1. Glossary.	
d	Branch duration, i.e., the time (and chronostratigraphic gap) separating the first appearance of a species (or the splitting time of a node) from the divergence linking it to the rest of the clade.
t	Time elapsed after a divergence over which a clade of any size might accrue new species, sampled specimens or become extinct.
T	Duration of interval with unique diversification and/or sampling rates.
S	Sum of species durations within a clade, usually given in lineage-million-years (Lmyr).
λ_i	Cladogenesis (origination) rate in interval-slice i .
μ_i	Extinction rate in interval-slice i .
ψ_i	Sampling rate in interval-slice i .
Φ_i	Probability of ever sampling a species and/or any successors (= a clade of unknown size) when only one species is present at the outset of time-slice i .
Φ'_i	Probability of ever sampling a species and/or any successors within a particular interval when only one species is present at the outset of interval-slice i .
v_i	Probability of sampling a species and/or any successors <i>within</i> interval-slice i .
ζ_i	Probability of sampling a species and/or any successors <i>after</i> interval-slice i when only one species is present at the outset of time-slice i .
$E[d_i]$	Expected branch duration for a species with first appearance in interval-slice i given origination, extinction and sampling rates. This is also the expectation for an unsampled ancestor with descendants that diverged in interval-slice i .

1124

Table 2. The probability of sampling a clade of unknown size given origination and extinction = 0.4 and different levels of sampling (ψ). Φ_B gives the global estimate for a clade given Bapst (2013). Φ_D gives the global estimate given Didier et al (2017). med. Φ_i gives the median 100-kyr estimate that allows for variation in diversification and sampling.			
ψ	Φ_B	Φ_D	med. Φ_i
0.400	0.6180	0.6180	0.6259
0.126	0.4260	0.4260	0.4297
0.040	0.2702	0.2701	0.2716
0.013	0.1628	0.1627	0.1618
0.004	0.0955	0.0951	0.0867

1125