Chapter 2.

Fossil Localities of the Baynunah Formation

Faysal Bibi, Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science,
Invalidenstrasse 43, Berlin 10115, Germany. faysal.bibi@mfn-berlin.de

Mark Beech, Historic Environment Department, Abu Dhabi Department of Tourism & Culture, PO
Box 2380, Abu Dhabi, United Arab Emirates

Andrew Hill, Department of Anthropology, Yale University, PO Box 208277, New Haven, CT 06520,
USA

Brian Kraatz, Department of Anatomy, College of Osteopathic Medicine of the Pacific, Western
University of Health Sciences, 309 E. Second Street, Pomona, CA 91766-1854, USA

Abstract

Fossil localities of the Baynunah Formation are here described, including previously known localities and those discovered by fieldwork since 2002. Site descriptions and coordinates are given, including updated coordinates for previously known localities. Many localities have been damaged or entirely lost to development activities.

Running head: *Fossil localities*

Introduction

The history of paleontological work in the Al Dhafra (formerly Al Gharbia or Baynunah) region of western Abu Dhabi Emirate is given by Beech et al. (this volume) and was presented in detail by Hill et al. (1999) and later by Hill et al. (2012) and Bibi et al. (2013), and only a summary is given here. Vertebrate fossils were first documented in oil survey reports starting in the late 1940s. In 1979 and 1981, fossils were collected by Peter Whybrow of the Natural History Museum in London (NHM), and in 1983-1984 by Walid Yasin of the Al Ain Museum and a German archaeological team. The latter collection included the hippo mandible AUH 457 (Boisserie et al. 2017; Boisserie and Bibi, this volume). The NHM-Yale expedition, led by Peter Whybrow and Andrew Hill and also involving Yasin, was active in the field between 1982-1994, with some specimens collected as late as 1998. Results of the NHM-Yale expedition were published in a 1999 volume (Whybrow and Hill 1999). In 2002-2004 sporadic missions led separately by Bibi under the Public Works Department, and Beech under the Abu Dhabi Islands Archaeological Survey resulted in the discovery of new sites and recovery of new fossil material. From 2006 onwards, an annual expedition co-directed by Bibi, Hill, and Beech, founded on an agreement between Yale University (where both Bibi and Hill were at the time) and the Abu Dhabi Authority for Culture and Heritage (ADACH; now Department of Culture and Tourism, DCT). Yasin was involved in fieldwork between 2006 and 2009.

Whybrow and Clements (1999) described the fossil localities of the Baynunah Formation as discovered by the NHM-Yale team, and provided GPS coordinates for many of these based on the UAE Nahrwan datum (Whybrow, pers. comm. to F.B. 2002), rather than the now ubiquitous WGS 84 datum. The coordinates in Whybrow & Clements (1999) also fall tens to hundreds of meters from known site points, such as concrete survey markers at SHU 6 or JBR 1, even after conversion to the WGS 84 datum. It is highly likely that the published coordinates were affected by Selective Availability, by which public GPS signals were intentionally degraded, and which was only discontinued in 2000.

We attempted to relocate the localities described by Whybrow & Clements as closely as possible, though this was not always straightforward. We used the same site names and locality numbers to the degree possible, creating new localities for fossiliferous areas that were not previously

documented. In 2006, Beech created an official paleontological database for ADACH (now DCT) and all locality abbreviations were updated to three-letter abbreviations (previously one or two-letter). In 2002, Bibi introduced a sub-locality system to further constrain collecting areas. For example, SHU 2-2 represents sub-locality 2 in locality SHU 2 on the island of Shuwaihat. We quickly realized that this was unnecessary as almost every specimen collected as of 2002 was associated with a GPS coordinate usually accurate to between 6 and 8 m. This made the exact delineation of site and locality boundaries — a difficult and fairly arbitrary practice to begin with — less critical because the actual position of each specimen in relation to every other – and to the local stratigraphy – could always be reconstructed or revisited. The designation of sub-localities was not made consistently after 2007.

Our field strategy from 2002-2014 was to collect and catalog all mammal specimens that were identifiable to the family level. Collection of common taxa, such as crocodiles, turtles, fish, and ratite eggshell occurred less consistently, typically restricted to either more complete specimens, or to specimens collected as vouchers to document new collecting areas. All specimens were given field numbers at the moment of collection, typically one field number per specimen, but some collections of multiple specimens from the same location share a single field number. These were written in field notes, entered into the catalog, and written on collection bags and jackets, but never directly on the specimen itself. Field numbers have the format YEAR-NUMBER, for example 09-26, whereby 09 represents the field season starting year – 2009 – and 26 is a the collection number. Field seasons normally ran from December into January of the following year, but the field number always references the year in which the field season began. In our example, specimens collected in January 2010 would still carry the 09 number if fieldwork was initiated in December of 2009. Normally, within a few days of collection, all specimens were given official AUH numbers by FB, who maintained the most up-to-date catalog. Specimens belonging to different individuals or taxa and collected under a single field number were given separate AUH numbers. An updated copy of the fossil catalog was submitted to the Abu Dhabi government agency (presently DCT) at the end of each field season.

The large majority of fossil specimens are in Abu Dhabi and under the care of the DCT. A small collection, mainly that from 1983-1984, as well as those from 2006, are at the Al Ain Museum.

Holotypes found by the NHM-Yale expedition are housed at the Natural History Museum, London, and a small collection of AUH-numbered and unnumbered specimens are currently still there, presumably pending return to Abu Dhabi. A small number of specimens collected by Whybrow prior to 1982 are also permanently housed at the NHM, and numbered with ‘M’ numbers, including the type mandible of *Archaeopotamus qeshta*, M49464 (Boisserie et al., 2017; Boisserie and Bibi, this volume). A single specimen, the gomphothere molar 1965 I 112 (Madden et al. 1982; Sanders, this volume; Tassy 1999), is housed at the Bayerische Staatssammlung für Paläontologie und Geologie in Munich, Germany. Unfortunately, specimens collected under Bibi in 2002 and 2003 (AUH 790–1049), were taken to a private collection in Abu Dhabi. Attempts to return these to the public authority of the Department of Culture and Tourism are ongoing. We have tried to include these fossils in this volume to the degree possible, relying on photographs, notes, and measurements taken by FB before these specimens became inaccessible in 2004. Notable finds include the partial skeleton of a giraffe (Bibi, this volume), and the maxillary palate of a gomphothere proboscidean (Sanders, this volume).

FIGURE 2.1 NEAR HERE. FULL PAGE WIDTH.

The Baynunah fossil localities are scattered across the Al Dhafra region, mostly within proximity to highway E11, which runs between Abu Dhabi and Saudi Arabia (Fig. 2.1). This was previously known as the Abu Dhabi-As Sila highway, and west of Ruwais its course was recently (2013-2014) detoured to a newly-built segment located to the south. The old segment, running between Ruwais and Shuwaihat, is now labelled on maps as Al Rubban St.

The localities themselves are spread out over an east-west distance of more than 200 km, and a north-south distance of 25-40 km (covering an area about the size of Lebanon or Jamaica). Prior to increased development work in the region, access to fossil sites was generally easy as many dirt roads cut through the area, and driving off-road was possible on firm ground (except in the coastal sabkha areas). The current phase of exploration since 2002 has collected many more fossils from known sites, and recorded many new sites, in particular in areas inland from the coast. Much of this inland region between the coast and the sand deserts to the south is comprised of low-relief landscapes with

extensive and highly reflective and deflated carbonate surfaces of the upper Baynunah Formation. These carbonates are generally of algal-mat origin (Bibi et al., 2012; Schuster, this volume) and are often rich in ostracods and gastropod (cerithid) molds. They match the lithology, composition, and general elevation of carbonates in the upper parts of coastal sections such as at Hamra, Kihal, and Jebel Barakah. These carbonates are not ideal for bone preservation, but they do record vertebrate trackways such as the proboscidean trackway sites at Mleisa and Bida' Al Moutawi'a (Bibi et al., 2012; Bibi et al., this volume). Occurrences of vertebrate fossil remains deep inland are few, but notable occurrences are at Jaw al Dibsa and Matlaiyyat.

The mapping of all fossil localities and areas within Abu Dhabi Emirate is shared with the Abu Dhabi government GIS system via the Abu Dhabi Systems Information Centre (ADSIC) GeoPortal. The Department of Culture and Tourism participates in the electronic No Objection Certificate (e-NOC) and eMePs systems that control and regulate development within Abu Dhabi Emirate. Developers are not permitted to build on or destroy major cultural heritage sites, including fossil localities, without going through these permitting controls. This system has helped to preserve many of the existing fossil localities, as well as leading to the discovery of new fossil localities (e.g. at Ruwais). Nonetheless, over the years many localities have become inaccessible to scientific investigation due to fences, pipelines, highway and railway construction, or appropriation of the land for military or industrial use. Between 2003 and 2014, we documented disturbance or difficulty of access at multiple sites (e.g. Shuwaihat, Hamra, Ras al Qal'a, Ruwais, Mleisa), including Jebel Barakah, the stratotype of the Baynunah Formation (Whybrow, 1989). Some efforts have been made by ADACH/DCT to fence in some localities to deter vehicular traffic or other disturbances (e.g. BIM4, DBS3, MLS1, MLS2, RUM 7, UAI). But for the most part there has been little success in protecting most coastal sites from development, particularly rich and emblematic localities at Shuwaihat and Hamra (Figs. 2.1, 2.2).

What follows is a description of all localities from which fossil remains have been documented, including those listed by Whybrow & Clements (1999) as well as new localities discovered since 2002. All fossiliferous localities are documented in Table 1. All localities are given three-letter abbreviations. Old one or two-letter abbreviations are indicated as applicable. English

transliteration of Arabic place names is generally an inconsistent practice, and here we have followed spellings as taken from local maps, road signs, or as previously used. We have attempted to remain as consistent as possible. Arabic place names follow local spelling, when known.

FIGURE 2.2 NEAR HERE. 1.5 COLUMN WIDTH

Localities of the Baynunah Formation

Baynunah North بينونة شمال

Abbreviation: BYN

Description: Small hills inside municipal plantation grounds on the north side of the Abu Dhabi-As Sila E11 highway. Localities BYN 1 to 3 were designated by Beech while surveying for the projected route of the Etihad Railway. BYN 1 was the most productive site. BYN 2 is a point location within BYN 1 and is not a fossil site.

Bida' al-Ghinia بدع الغنية

Abbreviation: BGH

Description: An inland exposure of carbonates with a single locality preserving multiple proboscidean trackways, discovered by local ranger Mubarak bin Hazeem and studied by our team in December 2017. This marks the easternmost locality from which trackways have been documented to date.

Bida' Al Mutawi'a بدع المطاوعة

Abbreviation: BIM

Description: These localities lie about 35 km south of the coast from Jebel Baraka, near the town of Bida' Al Mutawi'a. BIM 1-3 are Late Neolithic archaeological localities (Crombe 2000) and are not considered here. BIM 4 is a proboscidean trackway site that was fenced in for protection. Fossil eggshell and fossil wood were collected from BIM 5, but exact location of these is not certain, and may be some distance from BIM 4.

Bin Jawabi بن جوابي

Abbreviation: BJW (previously BJ)

Description: Whybrow & Clements (1999) provided no information about this locality except that a few fragments were collected from here. The only cataloged specimens are AUH 637 to 640, all collected in 1994. The exact location of this locality is unknown.

Gerain al-Aysh قرين العيش

Abbreviation: GAA (previously R)

Description: An extensive area of fossiliferous outcrops, today located within the grounds of a palace at the time belonging to Sheikh Sultan bin Zayed. This is the same as Ras al Aysh in Whybrow & Clements (1999). The NHM-Yale team collected here in 1991 and 1992, before the area became palace grounds. Our team was granted entry for some days in 2008 and 2010, during which a large number of specimens were collected, including a well-preserved crocodile cranium from GAA 4 (AUH 1568).

Ghayathi غياثي

Abbreviation: GHA (previously G)

Description: Whybrow & Clements (1999) describe this locality as a small quarry near the town of Ghayathi from which some bivalve fossils were collected (AUH 335). It is also stated as being located “about 18 km north, and about 1 km east of the main road from Ghayathi” but the GPS coordinates provided by Whybrow & Clements (Nahrwan datum) indicate a location close to the highway and just north of the town, in the vicinity of which several small outcrops of Baynunah Formation are present. This site was not visited by our team and the exact collection location is unknown.

Al-Ghuddah

Abbreviation: GUD (previously GU)

Description: GUD 1 is described by Whybrow & Clements (1999) as a jebel (hill) located south of the highway, and at which bone fragments and ratite eggshell were found. A gastropod is also cataloged

from here. Maps and satellite imagery indicate two jebels at a location matching this name and description (approximate coordinates given in Table 1). This site was not visited by our team and the exact collection location is unknown.

Hadwaniya الحضوانية

Abbreviation: HAD

Description: An extensive area of exposures just to the west of Gerain al-Aysh, quite rich in fossil material. When surveyed in January 2011, a large quarry had been removing sediment from Baynunah Formation outcrops at HAD 1 (Fig. 2.3).

FIGURE 2.3 NEAR HERE. FULL PAGE WIDTH.

Hadwaniyya South الحضوانية جنوب

Abbreviation: HDS

Description: Low exposures south of the old Abu Dhabi-As Sila highway, across from Gerain al-Aysh. Editor BK suffered a stroke here on the day of discovery of the site, 14 Jan. 2011 (he recovered).

Hamra الحمرا

Abbreviation: HMR (previously H)

Description: A large and productive area of thick exposures (Fig. 2.2). Nine localities were described in Whybrow & Clements (1999), though fossils are cataloged from only HMR 1 to HMR 6. HMR 1, 2, 4, 5, and 6 are located north of the old Abu Dhabi-As Sila highway, while HMR 3 is ~1.5 km south of the highway. The most productive localities are HMR 1, located at the northern end of the exposures by the coast, and HMR 5, comprising a large jebel and surrounding areas located just north of the old Abu Dhabi-As Sila highway. HMR 6 was described as 'near to the top of the jebel' that we understand is in the same area as HMR 5. Though fossils collected by the Whybrow-Hill expedition were assigned to either HMR 5 or 6, we were unable to distinguish the two collecting areas and assigned all newly collected material from the area as from HMR 5. HMR 7 is described as north of the old

highway, while HMR 8 and 9 are described as survey points (Whybrow & Clements 1999). No fossils are cataloged from any of these three localities.

Harmiyah الهرمية

Abbreviation: HAR (previously Y)

Description: These are low exposures just north and south of the old Abu Dhabi-As Sila highway.

Most specimens are from HAR 1, with only a handful from HAR 2 through 4. Also spelled as Al Harmia. A new exit road from the E11 highway heading north was recently constructed, partly damaging this locality.

Jaw al-Dibsa جو الدبسا

Abbreviation: DBS

Description: This is a fairly featureless area of low relief, situated largely within a municipal plantation area approximately 20 km south of the coast. A few weathered bones were collected from the surface at DBS 1. DBS 2 is a highly deflated surface from which a partial proboscidean mandible and other fragments were excavated in December 2008 (AUH 1282, Fig. 2.4). DBS 3 has abundant fossil wood (e.g. AUH 1077).

FIGURE 2.4 NEAR HERE. 1.5 COLUMN WIDTH.

Jebel Barakah جبل براكه

Abbreviation: JBR, previously B

Description: This site is located on the coast and is the type section of the Baynunah Formation (Whybrow 1989), including also exposures of the Shuwaihat Formation at its base. There are five designated localities at Jebel Barakah, none of which are particularly rich. Localities JBR 1 to 4 are located on or around the main jebel and sea cliffs below it. JBR 5 covers most of the low exposures to the west and southwest of the jebel. Notable finds include the type mandible of *Archaeopotamus geshta* at JBR 2 (M49464, Boisserie et al. 2017; Boisserie and Bibi this volume). The area was already

the site of occasional military activity by the early 2000s, and became increasingly difficult to access over the years. By 2011 the site was encircled by a fence and the area was designated off limits.

Kihal الكحال

Abbreviation: KIH (previously K)

Description: An area of exposures north of the E11 highway and reachable by a dirt road leading to Thumayriyyah Island. The most productive areas are the east-facing outcrops to the east and north of the main jebel (Fig. 2.5). Whybrow & Clements (1999) described these as sites K1 and K2, though from their descriptions it appears that K2 refers to a trigonometric point on top of the jebel located within the K1 area. All Yale-NHM fossils collected from Kihal were cataloged as from K1, while fossils we collected from the same outcrops were assigned to KIH 2, and these localities are to be considered synonymous. Notable finds from this locality include AUH 1134, a ratite pelvis (Louchart et al. this volume). None of the other Kihal localities were very productive. KIH 3 refers to the jebel south of KIH 1 and just north of the highway. KIH 4 and KIH 5 refer to small, low relief collecting areas to the north of KIH 1 and closer to the coast.

FIGURE 2.5 NEAR HERE. 1.5 COLUMN WIDTH.

Jebel Dhanna جبل الظنة

Abbreviation: JDH (previously JD)

Description: Jebel Dhanna is a low mountain rising about 90m above sea level, comprised of old deformed sediments (probably the Mesozoic Hajjar Group) that have been uplifted by a salt diapir (Glennie 2001). Most of the jebel is part of a petroleum facility and is off limits. All localities here were named by the NHM-Yale expedition (Whybrow & Clements 1999). JDH 1 is a small outcrop close to the road, located to the east of Jebel Dhanna. There is no JDH 2. JDH 3, 4, and 5 are very close to each other and mark a set of outcrops by the ferry terminal to the northwest. These have been heavily modified by further development of the terminal and nearby construction, but our team was still able to find some fossils at JDH 4 in December 2006. A few specimens from the NHM-Yale

expeditions come from JDH6, but no information was provided on this locality by Whybrow & Clements (1999) and its exact location is not known.

Jebel Mimiya جبل المميا

Abbreviation: MIM (previously M)

Description: Just east of the town of Al Mirfa, quarried faces of two low hills provide some of the clearest exposures of Baynunah sediments at any site (MIM 1, Fig. 2.6; Friend 1999). The NHM-Yale expedition collected quite a few specimens here, but our team only found a total of three. The best of these was a juvenile proboscidean mandible collected in 2014 (AUH 1779; Sanders 2017; Sanders this volume). In December 2017, exploration of another outcrop ~300 m to the west (MIM 2) revealed what may be highly eroded proboscidean footprints, potentially making this the only fossil trackway known from the coastal Baynunah sites (Bibi et al. this volume, fig. 17.5).

FIGURE 2.6 NEAR HERE. ONE COLUMN WIDTH.

Matla'iyat المطلعيات

Abbreviation: MTL

Description: This is a low-relief area located ~30 km inland and around 20 km west of Umm al Ishtan. A few fossils were collected from here, found on or emerging from highly deflated surfaces.

Mleisa المليسة

Abbreviation: MLS

Description: Extensive surface exposures of carbonates preserving proboscidean and ungulate trackways, first brought to the attention of M. Beech by local ranger Mubarak bin Rashid Al Mansouri in 2001 (Fig. 2.7). MLS 1 marks the location of a proboscidean herd, while MLS 2 contains isolated proboscidean prints next to an ungulate trackway (Bibi et al., 2012; Bibi et al., this volume). The localities themselves are protected by fences, and access is restricted due to oil pipeline extensions, railway construction, and the presence of the Houbara Conservation Protected Area.

FIGURE 2.7 NEAR HERE. 1.5 COLUMN WIDTH.

Niqa

Abbreviation: NIQ

Description: NIQ 3 is a surface exposure of carbonate located about 15 km northeast of Ghayathi and preserving what may be one or more prints of a hippopotamid (Bibi et al. this volume; Higgs et al. 2005). NIQ 1 and 2 are archaeological collection points, not fossil sites.

Ras al Qal'a راس القلعة

Abbreviation: RAQ (previously Q)

Description: A large area of outcrops on the coast, to the northwest of Kihal. A good dirt causeway leading to Thumayriyah island diverges leads west over the sabkha to the exposures. Most fossils come from locality RAQ 2. This general area has seen much bulldozing and earth movement activity in recent years. Also known as Qila' or Jila' (القلاع).

Ras Dubai'ah راس ضبيعة

Abbreviation: RDB (previously R)

Description: Ras Dubai'ah is a fairly large collecting traversing both sides of the E11 highway between Kihal and Ruwais. Whybrow & Clements (1999) wrote that a fragment of a rhinocerotid molar was discovered here, but that otherwise the Ras Dubai'ah area was never surveyed and had become a restricted area. The specimen catalog inherited from Whybrow lists specimens as coming from localities R1 and R2 (as well as one from R3, but we believe this to be a typo). We are not sure of the exact position of these localities. We identified RDB 1 (R1) based on the recollection of A. Hill and W. Yasin in the field (pers. comm. to FB). We are uncertain of the location of R2, but it may have been close to RDB 1. We designated new localities RDB 3, approximately 5 km to the east, and RDB 4, near a plantation site south of the highway from RDB 3. RDB 4 was initially (2002-2003) noted in Bibi's field notes as 'Municipality' locality (MU 1).

Rumaitha الرميثة

Abbreviation: RUM

Description: The Rumaitha site, in particular locality RUM 7, is among the easternmost points from which Baynunah fossils have been recovered. More recently the British Geological Survey identified some Baynunah outcrops even further east, close to the Al Dhafra military base. Some fossil fragments were found and logged at RUM by Beech in 2002 (Beech 2005), but no specimens have been cataloged from this site.

Ruwais الرويس

Abbreviation: RUW

Description: Ruwais Central and North (RUW C, RUW N) were surveyed as part of a salvage project in 2002 led by Beech. TAKREER, the Abu Dhabi Oil Refining Company (part of the Abu Dhabi National Oil Company, ADNOC) commissioned an environmental impact assessment of an area close to Ruwais where the BeAAT Central Environment Protection Facilities were to be built. This was located east of the Ghayathi Road (highway E15), not far from its junction with Ruwais. Just to the southeast is RUW SE, a site discovered in 2012 by an environmental impact assessment carried out for the route of the Etihad Railway. Our team briefly visited RUW SW in 2014, collecting numerous specimens including the holotype of a newly named rodent (AUH 1750, Kraatz this volume).

Shahriya الشهريّة

Abbreviation: SHA

Description: Exposed carbonate surface preserving highly weathered proboscidean trackways. Located within a protected wilderness conservation area, it was first discovered by Ahmad El Hajj and Jaber el Merri in 2007 and visited by our team in December 2017.

Shuwaihat الشويّهات

Abbreviation: SHU (previously S)

Description: Shuwaihat is an island connected to the mainland by a raised causeway (Figs. 2.2 and 2.8). There, the exposures of the Baynunah and Shuwaihat Formations are at their most typical and beautiful. This is one of the most intensely collected areas of the Baynunah Formation. The NHM-Yale expedition named five localities which we re-identified as closely as possible. SHU 1 is currently within a military base on the eastern side of the island and was never accessible to us. SHU 2 is an isolated jebel with surrounding outcrops to the northwest of the island. Subsite SHU 2-2 was designated by our team and targeted over many years for microvertebrate remains, notably producing rodents, squamates, and a monkey tooth (Kraatz et al. 2013; Kraatz, this volume; Head and Müller, this volume; Gilbert and Hill, this volume, Gilbert et al. 2014). In contrast, the majority of microvertebrates recovered by the NHM-Yale expedition came from localities SHU 1 and SHU 4 (de Bruijn and Whybrow 1994, Bruijn 1999). SHU 1 was never accessible to us, and efforts to recover microvertebrates at SHU 4 were not successful.

SHU 3 is a large jebel with surrounding outcrops to the southwest of the island. Whybrow & Clements (1999) noted this was known locally as Jebel Mershed, and though they referred to it as the westernmost jebel, it is in fact the southernmost (S2 is the westernmost). SHU 4 is the northeastern flank of the island's main western jebel. SHU 5 is an old oil well, and not a fossil site. SHU 6 is the excavation from which a proboscidean skeleton was recovered (Andrews 1999), and which was marked with a concrete datum by the NHM team in 1994.

We also named several additional localities. SHU 7 is the southwestern, southern, and southeastern slopes of the island's main western jebel. The largest east-facing gully demarcates this site from SHU 4. A thick covering of surface sand marks the boundary between SHU 7 and SHU 8, which covers the northwestern flank of the jebel. Neither of these areas was very fossiliferous. SHU 9 and SHU 10 cover the northern and southern ends, respectively, of the same jebel. SHU 11 was named for some fossil material recovered from Zabbut, the tiny island about 1.2 km north of Shuwaihat, which can be accessed either by boat or, at low tide, by car. SHU 12 is a jebel to the northwest of Shuwaihat, located between SHU 2 and SHU 8.

FIGURE 2.8 NEAR HERE. 1.5 COLUMN WIDTH.

Sir Bani Yas صير بني ياس

Abbreviation: SBY

Description: The island of Sir Bani Yas (now a wildlife park) was visited in December 2014.

Outcropping there along the edges of older strata uplifted by salt diapir activity are sedimentary sequences that appear to belong to the upper Baynunah Formation. No vertebrate remains were recovered, but ostracods were found in carbonates at the northern end of the island (SBY 1; Mazzini & Kovacova this volume).

Talfaha طلفحا

Abbreviation: TAL

Description: A small set of low-lying hills ~3km southeast of Jebel Barakah. The site was mistakenly spelled "Talahafa" in Bibi et al. (2006). Only aepeornithid-type fossil eggshells have been collected from here.

Tarif الطريف

Abbreviation: TAR (previously T)

Description: Locality TAR 2 produced a partial cranium of the bovid *Tragoportax* in 1992 (AUH 442, Gentry 1999; Bibi, this volume; see also Hill et al. 1999). Localities T1 and T3 are survey points, not fossil localities. We have not relocated these localities and their status is undetermined. The GPS coordinates provided by Whybrow & Clements (1999) suggest they all lie south of the highway and close to the Habshan road.

Thumayriyah الثميرية

Abbreviation: THM (previously TH)

Description: Thumayriyah is a small coastal island reached by a raised causeway leading north from the E11 highway. At the northwestern end of the island are Baynunah exposures as well as a private

house that was abandoned when we surveyed there in 2010 and 2011. The entire island is considered a single collecting area (THM 1), though all our team's finds come from the small peninsula at its northern end.

Umm al Ishtan أم العشتان

Abbreviation: UAI

Description: A deflated, low-relief area southeast of Jaw al Dibsa. UAI 1 is an area with abundant fossil wood that was fenced off for protection by ADACH. UAI 2 is a point location about 500m to the northwest, by the road near a ranger station, at which fossil wood was noted. UAI 3 is located within, and considered synonymous with, UAI 1. UAI 4 lies to the northwest. A proboscidean pelvis, right femur, and tibia were excavated here in February 2006 by Beech. The left femur of the same individual was recovered in 2014 (AUH 1740, Sanders this volume).

Umm al Kabir أم الكبير

Abbreviation: UAK

Description: An area northwest of Mleisa and east of the Ghayathi road. UAK 1 is a large deflated surface exposure of indurated carbonate-rich green silt, first discovered by Steve Raynor and the Emirates Natural History group on a trip to visit the Mleisa 1 site in December 2006. Multiple locations preserve what appear to be pedestaled proboscidean footprints (see Bibi et al. this volume, fig 17.11), along with eggshell, enamel, and crocodile fragments (AUH 1690-1692). UAK 2 is another area with similarly raised mounds that appear to be proboscidean footprints. North of UAK, a large exposure of Baynunah sediments is overlain by carbonate-cemented aeolian dunes (Fig. 2.9), most likely formed during Pleistocene glacial maxima (Glennie and Singhvi 2002; Schuster this volume).

FIGURE 2.9 NEAR HERE. FULL PAGE WIDTH.

Acknowledgements

The main expeditions for the work described here (2006-2014) took place under an agreement between Yale University at the Abu Dhabi Authority for Culture and Heritage (now the Department of Culture and Tourism, DCT). Main project support has come from ADACH / DCT, the Yale Peabody Museum of Natural History and Yale University, the National Science Foundation (grant OISE-0852975 to Bibi), a Leibniz-DAAD scholarship (to Bibi), the Revealing Hominid Origins Initiative (NSF 0321893 to T. White and F. C. Howell), and the Institut de Paléoprimatologie, Paléontologie Humaine: Évolution et Paléoenvironnements (iPHEP) at the University of Poitiers. Bibi also acknowledges support from a 2010 Fine Outreach for Science GigaPan Fellowship (Carnegie Mellon University). We thank Patricia Holroyd, Nancy Stevens, and Tim White for suggestions that helped improve this chapter.

References

- Andrews P. (1999) Taphonomy of the Shuwaihat proboscidean, late Miocene, Emirate of Abu Dhabi, United Arab Emirates. In: Whybrow P. J. Hill A. P. (ed) Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates, Yale University Press, New Haven, p 338-353
- Beech M. (2005) The late Miocene fossil site at Ruwais. In: Beech M. Hellyer P. (ed) Abu Dhabi 8 Million Years Ago: Late Miocene Fossils from the Western Region, Abu Dhabi Islands Archaeological Survey, Abu Dhabi, p 21-33
- Beech M., Kraatz B., & Bibi F. (this volume). Introduction. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Bibi, F. (this volume). Bovidae and Giraffidae from the Baynunah Formation. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Bibi F., Hill A., Beech M. & Yasin W. (2013) Late Miocene fossils from the Baynunah Formation, United Arab Emirates: Summary of a decade of new work. In: Wang X., Flynn L. J. Fortelius M. (ed) Fossil Mammals of Asia: Neogene Biostratigraphy and Chronology, Columbia Univ. Press, New York, p 583-594
- Bibi, F., Kraatz, B., Beech, M., & Hill., A. (this volume). Fossil Trackways of the Baynunah Formation. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Bibi, F., B. P. Kraatz, N. Craig, M. Beech, M. Schuster, and A. Hill. 2012. Early evidence for complex social structure in Proboscidea from a late Miocene trackway site in the United Arab Emirates. *Biology Letters* 8(4):670-673.
- Bibi F., Shabel A. B., Kraatz B. P. & Stidham T. A. (2006) New fossil ratite (Aves: Palaeognathae) eggshell discoveries from the Late Miocene Baynunah Formation of the United Arab Emirates, Arabian Peninsula. *Palaeontologia Electronica* 9:2A: 13p

- Boisserie, J.-R. & Bibi, F. (this volume). Hippopotamidae from the Baynunah Formation. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Boisserie J.-R., Schuster M., Beech M., Hill A. & Bibi F. (2017) A new species of hippopotamine (Cetartiodactyla, Hippopotamidae) from the late Miocene Baynunah Formation, Abu Dhabi, United Arab Emirates. *Palaeovertebrata* 40:doi: 10.18563/pv.41.1.e2
- Bruijn H. d. (1999) A late Miocene insectivore and rodent fauna from the Baynunah Formation, Emirate of Abu Dhabi, United Arab Emirates. In: Whybrow P. J. Hill A. P. (ed) *Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates*, Yale University Press, New Haven, p 186-197
- Crombe P. (2000) A neolithic site at Bida Al Mitawaa in Western Abu Dhabi (U.A.E.). *Arabian Archaeology and Epigraphy* 11:9-14
- De Bruijn H. & Whybrow P. J. (1994) A Late Miocene rodent fauna from the Baynunah Formation, Emirate of Abu Dhabi, United Arab Emirates. *Proceedings Koninklijke Nederlandse Akademie van Wetenschappen* 97:407-422
- Friend P. F. (1999) Rivers of the lower Baynunah Formation, Emirate of Abu Dhabi, United Arab Emirates. In: Whybrow P. J. Hill A. (ed) *Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates*, Yale University Press, New Haven, p 39-49
- Gentry A. W. (1999) Fossil pecorans from the Baynunah Formation, Emirate of Abu Dhabi, United Arab Emirates. In: Whybrow P. J. Hill A. (ed) *Fossil Vertebrates of Arabia*, Yale University Press, New Haven, p 290-316
- Gilbert C. C., Bibi F., Hill A. & Beech M. J. (2014) Early guenon from the late Miocene Baynunah Formation, Abu Dhabi, with implications for cercopithecoid biogeography and evolution. *Proceedings of the National Academy of Sciences* 111:10119-10124
- Glennie K. W. (2001) Evolution of The Emirates' Land Surface: an Introduction. In: Al Abed I. Hellyer P. (ed) *United Arab Emirates: A New Perspective*, Second. Trident Press, London, p 9-27

- Glennie K. W. & Singhvi A. K. (2002) Event stratigraphy, paleoenvironment and chronology of SE Arabian deserts. *Quaternary Science Reviews* 21:853-869
- Head, J. & Müller, J. (this volume). Amphibians and squamates from the Baynunah Formation. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Higgs, W., Gardner, A., & Beech, M. (2005). A fossil proboscidean trackway at Mleisa, Western Region of Abu Dhabi, United Arab Emirates. In P. Hellyer, & M. Ziolkowski (Eds.), *Emirates Heritage Vol. 1 – Proceedings of the 1st Annual Symposium on Recent Palaeontological and Archaeological Discoveries in the Emirates* (pp. 21-27). Al Ain: Zayed Centre for Heritage and History.
- Hill A., Bibi F., Beech M. & Al-Tikriti W. Y. (2012) Before archaeology: life and environments in the Miocene of Abu Dhabi. In: Potts D. Hellyer P. (ed) *Fifty Years of Emirates Archaeology*, Ministry of Culture, Youth and Community Development, Abu Dhabi, p 20-23
- Hill A., Whybrow P. J. & Yasin W. (1999) History of palaeontological research in the Western Region of the Emirate of Abu Dhabi, United Arab Emirates. In: Whybrow P. J. Hill A. P. (ed) *Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates*, Yale University Press, New Haven, p 15-23
- Kraatz, B. (this volume). Rodents from the Baynunah Formation. In F. Bibi, B. Kraatz, M. Beech, & A. Hill (eds.) *Sands of Time: Late Miocene Fossils from the Baynunah Formation, U.A.E.* (pp. xxx). Cham: Springer.
- Kraatz B. P., Bibi F., Hill A. & Beech M. (2013) A new fossil thryonomyid from the Late Miocene of the United Arab Emirates and the origin of African cane rats. *Naturwissenschaften* 100:437-449
- Madden C. T., Glennie K. W., Dehm R., Whitmore F. C., Schmidt R. J., Ferfoglia R. J. & Whybrow P. J. (1982) *Stegotetrabelodon* (Proboscidea, Gomphotheriidae) from the Miocene of Abu Dhabi United States Geological Survey, Jiddah
- Sanders W. J. (2017) Horizontal tooth displacement and premolar occurrence in elephants and other elephantiform proboscideans. *Historical Biology* 1-20

- Tassy P. (1999) Miocene elephantids (Mammalia) from the Emirate of Abu Dhabi, United Arab Emirates: Palaeobiogeographic implications. In: Whybrow P. J. Hill A. (ed) Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates, Yale University Press, New Haven, p 209-233
- Whybrow P. J. (1989) New stratotype; the Baynunah Formation (Late Miocene), United Arab Emirates: lithology and palaeontology. *Newsletters on Stratigraphy* 21:1-9
- Whybrow P. J. & Clements D. (1999) Late Miocene Baynunah Formation, Emirate of Abu Dhabi, United Arab Emirates: fauna, flora, and localities. In: Whybrow P. J. Hill A. P. (ed) Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates, Yale University Press, New Haven, p 317-333
- Whybrow P. J. & Hill A. (1999) Fossil Vertebrates of Arabia, with Emphasis on the Late Miocene Faunas, Geology, and Palaeoenvironments of the Emirate of Abu Dhabi, United Arab Emirates Yale University Press, New Haven

Table 2.1. Localities from which fossils have been collected or otherwise documented. All GPS coordinates taken using datum WGS84, including updated coordinates for localities described by Whybrow & Clements (1999).

Area	Locality	N	E
Baynunah North	BYN 1	24.10852	52.92186
	BYN 2		
	BYN 3	24.10688	52.92297
Bida' Al Moutawi'a	BIM 4	23.76827	52.55319
	BIM 5	23.88926	52.62613
Bida' al-Ghinia	BGH 1	23.89575	53.39819
Bin Jawabi	BJW	unknown	unknown
Gerain al-Aysh	GAA 1	24.086	53.217
	GAA 2	24.07319	53.22441
	GAA 3	24.09572	53.20691
	GAA 4	24.117	53.16691
	GAA 5	24.12458	53.18455

	GAA 6	24.13386	53.188
	GAA 7	24.15775	53.13338
	GAA 8	24.1438	53.14283
Ghayathi			
	GHA 1	unknown	unknown
Al Ghuddah			
	GUD 1	23.97500	52.53333
Hadwaniyya			
	HAD 1	24.13047	53.10025
	HAD 2	24.12077	53.10369
	HAD 3	24.11233	53.10413
	HAD 4	24.10341	53.10841
	HAD 5	24.08763	53.12663
	HAD 6	24.09158	53.12183
	HAD 7	24.09627	53.12352
Hadwaniya South			
	HDS 1	24.04983	53.205
Hamra			
	HMR 1	24.11466	52.52563
	HMR 1	24.11619	52.52419
	HMR 2	24.10808	53.52405

	HMR 3	24.07544	52.53013
	HMR 4	24.10263	52.51755
	HMR 5	24.10313	52.52827
	HMR 6	24.10227	52.5245
Harmiyah			
	HAR 1	24.07047	53.34133
	HAR 2	24.07872	53.32658
	HAR 3	24.07421	53.32798
	HAR 4	24.05758	53.30555
Jaw Al Dibsa			
	DBS 1	23.82233	52.32616
	DBS 2	23.81436	52.31086
	DBS 3	23.83083	52.33143
Jebel Barakah			
	JBR 1	24.00894	52.33086
	JBR 2	24.00547	52.32758
	JBR 3	24.1013	52.52736
	JBR 4	24.00655	52.32702
	JBR 5	24.00197	52.33041
Jebel Dhanna			
	JDH 1	24.1466	52.63517

	JDH 2	na	na
	JDH 3	24.1773	52.57402
	JDH 4	24.18063	52.57177
	JDH 5	24.17308	52.57727
	JDH 6	unknown	unknown
Jebel Mimiya			
	MIM 1	24.08483	53.43519
	MIM 2	24.08602	53.43302
Kihal			
	KIH 1	same as KIH 2	
	KIH 2	24.11891	53.02036
	KIH 3	24.10402	53.02366
	KIH 4	24.13172	53.00638
	KIH 5	24.12508	53.00511
Matla'iyat			
	MTL 1	23.70497	52.27238
	MTL 2	23.70533	52.26319
Mleisa			
	MLS 1	23.9480	53.06141
	MLS 2	23.94433	53.14625
Niqa			

	NIQ 3	23.95248	52.918599
Ras al Qal'a			
	RAQ 1	24.15227	52.97711
	RAQ 2	24.15063	52.98319
	RAQ 3	24.14855	52.98019
	RAQ 4	24.15036	52.98483
Ras Dubay'ah			
	RDB 1	24.1263	52.84169
	RDB 2	unknown	unknown
	RDB 3	24.11886	52.88991
	RDB 4	24.08333	52.9
Rumaitha			
	RUM 11	24.02548	54.12572
Ruwais Central			
	RUW C	24.08391	52.75691
Ruwais North			
	RUW N	24.10833	52.76666
Ruwais South East			
	RUW SE	24.07497	52.78947
Shahriyya			
	SHA 1	23.83777	52.61447

Shuwaihat

SHU 1	24.11605	52.44405
SHU 2	24.11527	52.43222
SHU 3	24.10288	52.43925
SHU 4	24.11241	52.43761
SHU 5	24.12033	52.44038
SHU 6	24.11388	52.43788
SHU 7	24.10952	52.43769
SHU 8	24.11147	52.43641
SHU 9	24.11947	52.44313
SHU 10	24.11469	52.44616
SHU 11	24.13308	52.4355
SHU 12	24.11358	52.43291

Sir Bani Yas

SBY 1	24.356722	52.588917
-------	-----------	-----------

Talfaha

TAL 1	23.99394	52.3538
-------	----------	---------

Tarif (approximate location)

TAR 2	23.94927	53.69244
-------	----------	----------

Thumayriyah

THM 1	24.16126	53.0092
-------	----------	---------

Umm al Ishtan

UAI 1	23.69232	52.43328
UAI 2	23.69928	52.4223
UAI 3	23.69189	52.4353
UAI 4	23.73047	52.35672

Umm al Kabir

UAK 1	24.01138	52.86769
UAK 2	23.99512	52.85742

Figure Captions

Figure 2.1. Fossil localities of the Baynunah Formation (top) and collection intensity by number of specimens (below). By far the most productive localities, and among those the most threatened by development, are at Hamra and Shuwaihat. Stippled rectangles denote areas shown in Figure 2.

Figure 2.2. Specimen collection mapping at Shuwaihat (top left), Hamra (top right) and several eastern coastal localities (below). Site locations and boundaries are shown, and each translucent black dot represents a single collected fossil specimen (exact locations of specimens collected prior to 2002 are approximated).

Figure 2.3. Kite aerial photographs of Baynunah exposures at Hadwaniyah (HAD) 1. To the left, the main outcrop. Note the 3 team members prospecting on its slopes, as well as the recreational vehicle tracks above, and the dirt road and fence in the background. To the right, zoomed out photomosaic of the same site, showing major quarrying for sand in the foreground. Kite images by N. Craig. 3 January 2011.

Figure 2.4. A pile of fossil proboscidean bones forms an erosion-resistant mound on an otherwise old and deflated surface of Baynunah Formation sediments at Jaw Al Dibsa (DBS) 2. AUH 1282, 18 December 2008.

Figure 2.5. Baynunah exposures looking north at Kihal (KIH 1 and 2). A crouching Andrew Hill for scale. 1 January 2009.

Figure 2.6. Baynunah Formation outcrop at Jebel Mimyah (MIM) 1, a fossil site that has also been quarried for sand. 18 December 2014.

Figure 2.7. Typical inland landscape resulting from extensive exposure of hard white carbonate beds of the upper part of the Baynunah Formation. Near Mleisa (MLS) 2, 9 December 2006.

Figure 2.8. Looking south from Shuwaihat (SHU) 2, showing the typical flat-topped jebels of Baynunah Formation coastal exposures. Note the team members to the bottom right. The cars below them are parked on the aeolian beds of the underlying Shuwaihat Formation. In the middle distance SHU 12, 7, and 8, and the isolated jebel in the back center of the frame is SHU 3. 26 December 2008.

A high-resolution photomosaic (Gigapan) of this same view is viewable here:

<http://gigapan.com/gigapans/68192>

Figure 2.9. Photomosaic (Gigapan) of exposures north of Umm al Kabir (UAK) 1. Baynunah Formation sediments (e.g. red bed at left) are overlain by carbonate-rich Pleistocene dunes that formed during glacial intervals as well as more recent desert sands. F. Bibi and M. Schuster for scale to the right. 8 January 2011. Location: 24.047562°N, 52.836123°E. Also viewable at

<http://gigapan.com/gigapans/74057>

Fig 2.1

Fig 2.2

Fig 2.3

Fig 2.4

Fig 2.5

Fig 2.6

Fig 2.7

Fig 2.8

Fig 2.9